CHAPTER II.

PHYSIOGRAPHY.

§ 1. General Description of Australia.

- 1. Geographical Position.—(i) General. The Australian Commonwealth, which includes the island continent of Australia proper and the island of Tasmania, is situated in the Southern Hemisphere, and comprises in all an area of about 2,974,581 square miles, the mainland alone containing about 2,948,366 square miles. Bounded on the west and east by the Indian and Pacific Oceans respectively, it lies between longitudes 113° 9′ E. and 153° 39′ E., while its northern and southern limits are the parallels of latitude 10° 41′ S. and 39° 8′ S., or, including Tasmania, 43° 39′ S. On its north are the Timor and Arafura Seas and Torres Strait, on its south the Southern Ocean and Bass Strait. The extreme points are "Steep Point" on the west, "Cape Byron" on the east, "Cape York" on the north, "Wilson's Promontory" on the south, or, if Tasmania be included, "South East Cape."
- (ii) Tropical and Temperate Regions. Of the total area of Australia nearly 40 per cent. lies within the tropics. Assuming, as is usual, that the latitude of the Tropic of Capricorn is 23° 30′ S., its correct value for 1924 is 23° 26′ 57.02″, and it decreases about 0.47″ per annum, the areas within the tropical and temperate zones are approximately as follows:—

AUSTRALIA-AREAS OF TROPICAL AND TEMPERATE REGIONS.

(STATES AND TERRITORY PARTIALLY WITHIN TROPICS.)

· Areas.	Queensland.	Western Australia.	Northern Territory.	Total.
Within Tropical Zone Within Temperate Zone Ratio of Tropical part to whole State Ratio of Temperate part to whole State	Sq. miles. 359,000 311,500 0.535 0.465	Sq. miles. 364,000 611,920 0.373 0.627	Sq. miles. 426,320 97,300 0.814 0.186	Sq. miles. 1,149,320 1,020,720 0.530 0.470

Thus the tropical part is roughly about one-half (0.530) of the three territories mentioned above, or about five-thirteenths of the whole of Australia (0.386).

2. Area of Australia compared with Areas of other Countries.—It is not always realized that the area of Australia is nearly as great as that of the United States of America, that it is four-fifths of that of Canada, that it is over one-fifth of the area of the whole of the British Empire, that it is more than three-fourths of the whole area of Europe, and that it is about 25 times as large as Great Britain and Ireland or Italy. This great area, coupled with a limited population, renders the solution of the problem of Australian development a particularly difficult one. The areas of Australia and of other countries are given in the following table:—

AREA OF AUSTRALIA AND OF OTHER COUNTRIES.

Country.	Area.	. Country.	Area.
Continental Divisions—	Sq. miles.	Asia—	Sq. miles.
Europe	3,854,145	Russia	5,460,390
Asia	15,972,271	China and Dependencies	4,277,170
Africa	12,413,292	British India and Adminis-	, ,
North and Central America		tered Territories	1,094,300
and West Indies	8,604,072	Arabia	1,000,000
South America	7,316,764	Feudatory Indian States	711,032
Australasia and Polynesia	3,464,891	Persia	628,000
		Dutch East Indies	572,950
Total, exclusive of Arctic	F1 007 407	Turkey	483,656
and Antarctic Conts.	51,625,435	Japan (and Dependencies)	260,738
3		Afghanistan	245,000
Europe— Russia	1,680,730	Siam Iraq (Mesopotamia)	200,148 $143,250$
Russia France	212,659	Philippine Islands	115,026
Spain	194,800	Laos	96,500
Germany	182,213	Omán	82,000
Sweden	173,105	Bokhara	79,440
Poland	149,359	British Borneo, Brunei,	,
Finland	132,550	and Sarawak	77,106
Norway	124,964	Syria	60,000
Rumania	122,282	Cambodia	57,900
Italy and annexed Pro-	11-000	Nepál	54,000
vinces	117,982	Tonking	40,530
Serb, Croat, and Slovene	06 124	Annam	39,758
State	96,134	Federated Malay States Ceylon	27,506
Ireland	94,633	Ceylon Khiva	25,332 $24,310$
Lithuania	59,633	Malay Protectorate (inc.	24,010
Czecho-Slovakia	54,191	Johore)	23,486
Bulgaria	39,824	Cochin China	22,000
Iceland	39,709	Bhután	20,000
Hungary	35,921	Armenia	15,240
Portugal	35,490	Aden and Dependencies	9,000
Azerbaijan	33,970	Palestine	9,000
Greece	33,378	Timor, etc. (Portuguese In-	- 000
Austria	32,396	dian Archipelago)	7,330
Irish Free State	27,000	Cyprus	3,584
Georgia	25,760	Goa, Damao, and Diu	1,638
Latvia Spitzbergen, Bear and ad-	25,000	Straits Settlements	1,600 1,382
jacent islands	25,000	Kwantung	538
Albania	17,374	Hong Kong and Dependen-	000
Denmark	17,144	cies	391
Estonia	16,955	Wei-hai-wei	285
Switzerland	15,975	Bahrein Islands	250
Netherlands	13,205	French India (Pondicherry,	
Belgium	11,752	etc.)	196
Turkey	10,882	Kwang Chau Wan	190
Luxemburg	999	Maldive Islands	115
Danzig	754	Macao, etc	4
Andorra	191	Total Asia	15,972,271
Malta	118 65	Total, Asia	10,912,271
San Marino	38		
Monaco	8	l l	
Gibraltar	2	Africa-	
		French Sahara	1,544,000
Total, Europe	3,854,145	Anglo-Egyptian Sudan	1,014,400
		French Equatorial Africa	982,049

AREA OF AUSTRALIA AND OF OTHER COUNTRIES-continued.

Area.	Country.	Агеа.
Sq. miles.	Africa—continued.	Sq. miles,
909,654		•
617,600		790
484,800	St. Thomas and Principe	
473,089	Islands	360
440,000	Seychelles	156
		47
	Ascension	34
	Total, Africa	12,413,292
	i ·	
	No. 45 A A	
	Λ	0 = 00 00 -
275,000	TT. 14 3 Ct 4	3,729,665
94# 060	36 .	3,026,789
	A1 - 1	767,198
		590,884
223,600		100 504
222 180		162,734
		51,660
	*/\	48,290
	TT. 1	$46,740 \\ 44,275$
		44,164
		23,000
		19,332
	1 0 1 1	13,176
,	TT 1.4	10,204
80,000	TO 1/1 TT 3	8,598
74,112		-,
68,000	and Cayman Is	4,431
50,000	Bahamas	4,404
45,783	Porto Rico	3,435
42,460	Trinidad and Tobago	1,976
40,000	Leeward Islands	715
39,573	Guadeloupe and Dependen-	
31,000	cies	688
		516
		403
		385
		166
		132
11,716	1 1 1	93
10.010	Bermudas	19
	Watel N and C America	
	1 777 7 10	0.004.0=0
	and w. indies	8,604,072
	-	
	South Amorica	
	701	9 974 510
	A 12 '70 . 1.12	3,275,510
		1,153,119 $722,461$
	D-1: ·	514,155
795	Colombia (exc. of Panama)	014,100
	\$q. miles. 909,654 617,600 484,800 473,089 440,000 365,000 350,000 350,000 350,000 225,000 225,000 223,800 223,800 222,180 166,489 154,400 139,430 121,976 110,300 109,200 95,218 80,000 74,112 68,000 50,000 45,783 42,460 40,000 39,573	Sq. miles. 909,654 617,600 484,800 473,089 440,000 428,132 406,000 350,000 350,000 357,000 327,400 327,400 327,000 223,800 223,800 223,800 223,800 223,800 223,800 221,976 10,300 109,200 95,218 80,000 74,112 68,000 45,783 42,460 40,000 45,783 42,460 40,000 45,783 42,460 40,000 45,783 42,460 40,000 45,783 42,460 40,000 45,783 42,460 40,000 45,783 42,460 40,000 45,783 42,460 40,000 45,783 42,460 40,000 45,783 42,460 40,000 45,783 42,460 40,000 45,783 42,460 40,000 45,783 42,460 40,000 45,783 42,460 40,000 45,783 42,460 40,000 45,783 42,460 40,000 41,112 68,000 45,783 42,460 40,000 40,

Danish colony only. Total area has been estimated as between 827,000 and 850,000 square miles.

AREA OF AUSTRALIA AND OF OTHER COUNTRIES-continued.

Country.	Area.	Country.	Area.
South America—continued.	Sq. miles.	Australasia and Polynesia —continued.	Sq. miles.
Ch:1-	398,594	British Solomon Islands	11,000
7	289,828		11,000
D. Milah, Calana	174,155	New Caledonia and Depen-	0 540
	89,480	177:2:	8,548 7,083
Uruguay	72,153	ii *	
Paraguay Dutch Guiana	61,647	N. TT had a	6,449
D	54,291	French Establishments in	5,700
First Oriona	32,380	1 .	1 = 00
Falkland Islands and	32,000		1,520
Q(1 Q. a.m.)-	# e10	Territory of Western Samoa Gilbert and Ellice Islands	1,250
D O. nol Zono	$5,618 \\ 527$	Marianne, Caroline, and	1,011
Panama Canal Zone	321	1	960
Total, South America	7,316,764		385
Total, South America	7,510,704	11 0 0	389 210
		Samoa (U.S.A. part)	210 77
		1 37 (11 7 1 1 1	13
Australasia and Polynesia—		Norioik Island	10
Commonwealth of Australia	0.054.501	Martin American and	
D (I M O)	2,974,581	Total, Australasia and	9.464.001
New Zealand and Depen-	160,692	Polynesia	3,464,891
dencies	103,862	-	
Territory of New Guinea	91,000		
Papua	90,540	British Empire	13,406,145

The figures quoted in the table have, in most cases, been extracted from the Statesman's Year-Book for 1924.

3. Areas of Political Subdivisions.—As already stated, Australia consists of six States and the Northern and Federal Capital Territories. The areas of these, and their proportions of the total of Australia, are shown in the following table:—

AUSTRALIA-AREA OF STATES AND TERRITORIES.

State or Territor	State or Territory.			Percentage on Total.
		•	Sq. miles.	
New South Wales			309,432	10.40
Victoria			87,884	2.96
Queensland			670,500	22.54
South Australia			380,070	12.78
Western Australia			975,920	32.81
Tasmania			26,215	0.88
Northern Territory			523,620	17.60
Federal Capital Territor	У		940	0.03
m			2,974,581	100.00

- 4. Coastal Configuration.—(i) General. There are no striking features in the configuration of the coast; the most remarkable indentations are the Gulf of Carpentaria on the north, and the Great Australian Bight on the south. The Cape York Peninsula on the extreme north is the only other remarkable feature in the outline. In Year Book No. 1, an enumeration of the features of the coast-line of Australia was given (see pp. 60 to 68).
- (ii) Coast-line. The lengths of coast-line, exclusive of minor indentations, of each State and of the whole continent, and the area per mile of coast-line, are shown in the following table:—

AUSTRAL		LINE ANI	AREA TER MILL	THEREOF	•
State.	Coast-line.	Area ÷ Coast-line.	State.	Coast-line.	Area ÷ Coast-line.
	Miles.	Sq. miles.		Miles.	Sq. miles.
New South Wales(a)	700	443	South Australia	1,540	247
Victoria	680	129	Western Australia	4,350	224
Queensland	3,000	223	Continent (b)	11,310	261
Northern Territory	1,040	503	Tasmania	900	29
	1				

AUSTRALIA-COAST LINE AND AREA PER MILE THEREOF.

(a) Including Federal Capital Territory. (b) Area 2,948,366 square miles.

For the entire Commonwealth of Australia this gives a coast-line of 12,210 miles and an average of 244 square miles for one mile of coast-line. According to Strelbitski, Europe has only 75 square miles of area to each mile of coast-line, and, according to recent figures, England and Wales have only one-third of this, viz., 25 square miles.

- (iii) Historical Signifiance of Coastal Names. It is interesting to trace the voyages of some of the early navigators by the names bestowed by them on various coastal features—thus Dutch names are found on various points of the Western Australian coast, in Nuyts' Archipelago, in the Northern Territory and in the Gulf of Carpentaria; Captain Cook can be followed along the coasts of New South Wales and Queensland; Flinders' track is easily recognised from Sydney southwards, as far as Cape Catastrophe, by the numerous Lincolnshire names bestowed by him; and the French navigators of the end of the eighteenth and the beginning of the nineteenth century have left their names all along the Western Australian, South Australian, and Tasmanian coasts.
- 5. Geographical Features of Australia.—In each of the earlier issues of this Year Book fairly complete information has been given concerning some special geographical element. The nature of this information and its position in the various Year Books can be readily ascertained on reference to the special index following the index to maps and graphs at the end of this work.
- 6. Fauna, Flora, Geology, and Seismology of Australia.—Special articles dealing with these features have appeared in previous Year Books, but limits of space naturally preclude their repetition in each volume. As pointed out in 5 supra, however, the nature and position of these articles can be readily ascertained from the special index.

§ 2. Climate and Meteorology of Australia.*

- 1. Introductory.—In preceding Year Books some account was given of the history of Australian meteorology, including reference to the development of magnetic observations and the equipment for the determination of various climatological records. (See Year Book No. 3, pp. 79, 80.) In Year Book No. 4, pp. 84 and 87, will be found a short sketch of the creation and organization of the Commonwealth Bureau of Meteorology, and a résumé of the subjects dealt with at the Meteorological Conference of 1907. Space will not permit of the inclusion of this matter in the present issue.
- 2. Meteorological Publications.—The following publications are issued daily from the Central Meteorological Bureau, viz.:—(i) Weather charts. (ii) Rainfall maps.

 $[\]mbox{\ ^{\circ}}$ Prepared from data supplied by the Commonwealth Meteorologist, H. A. Hunt, Esquire, F.R. Met. Soc.

(iii) Bulletins, Victorian and Interstate, showing pressure, temperature, wind, rain, cloud extent, and weather. Similar publications are also issued from the divisional offices in each of the State Capitals.

Commencing with January, 1910, the "Australian Monthly Weather Report," containing statistical records from representative selected stations, with rain maps and diagrams, etc., is being published. Complete rainfall and other climatological data are published in volumes of meteorological statistics for each State separately.

The first text book of Australian meteorology, "Climate and Weather of Australia," was published in 1913.

In addition, fifteen Bulletins of Climatology have been published, particulars of which are given in preceding issues of the Official Year Book (see No. 12, page 54).

3. General Description of Australia.—A considerable portion (0.530) of three divisions of Australia is north of the tropic of Capricorn—that is to say, within the States of Queensland and Western Australia, and the Northern Territory, no less than 1,149,320 square miles belong to the tropical zone, and 1,020,720 to the temperate zone. The whole area of Australia within the temperate zone, however, is 1,825,261 square miles; thus the tropical part is about 0.386, or about five-thirteenths of the whole, or the "temperate" region is half as large again as the "tropical" (more accurately 1.591). By reason of its insular geographical position, and the absence of striking physical features, Australia is, on the whole, less subject to extremes of weather than are regions of similar area in other parts of the globe; and latitude for latitude Australia is, on the whole, more temperate.

The altitudes of the surface of Australia range up to a little over 7,300 feet, hence its climate embraces a great many features, from the characteristically tropical to what is essentially alpine, a fact indicated in some measure by the name Australian Alps given to the southern portion of the great Dividing Range.

On the coast, the rainfall is often abundant and the atmosphere moist, but in some portions of the interior it is very limited, and the atmosphere dry. The distribution of forest, therefore, with its climatic influence, is very uneven. In the interior, in places, there are fine belts of trees, but there are large areas also which are treeless, and where the air is hot and parching in summer. Again, on the coast, even so far south as latitude 35°, the vegetation is tropical in its luxuriance, and to some extent also in character. Climatologically, therefore, Australia may be said to present a great variety of features.

4. Meteorological Divisions.—(i) General. The Commonwealth Meteorologist has divided Australia, for climatological and meteorological purposes, into five divisions. The boundaries between these may be thus defined:—(a) Between divisions I. and II., the boundary between South and Western Australia, viz., the 129th meridian of east longitude; (b) between divisions II. and III., a line starting at the Gulf of Carpentaria, along the Norman River to Normanton, thence a straight line to Wilcannia on the Darling River, New South Wales; (c) between divisions II. and IV., a line from Wilcannia along the Darling River to its junction with the Murray; (d) between divisions II. and V., a line from the junction of the Darling and Murray Rivers, along the latter to Encounter Bay; (e) between divisions III. and IV., a line starting at Wilcannia, along the Darling, Barwon, and Dumaresq Rivers to the Great Dividing Range, and along that range and along the watershed between the Clarence and Richmond Rivers to Evans Head on the east coast of Australia; (f) between divisions IV. and V., a line from the junction of the Darling and Murray Rivers along the latter to its junction with the Murrumbidgee, along the Murrumbidgee to the Tumut River, and along the Tumut River to Tumut, thence a straight line to Cape Howe; (g) Tasmania is included in division V.

The population included within these boundaries at the Census of the 4th April, 1921, was approximately as follows:—

Division	I.	II.	III.	IV.	v.
Population	332,000	500,000	824,000	1,915,000	1,866,000

In these divisions, the order in which the capitals occur is as follows:—(a) Perth, (b) Adelaide, (c) Brisbane, (d) Sydney, (e) Melbourne, and (f) Hobart; and the climatological and meteorological statistics relating to the capital cities are dealt with herein in accordance with that order.

(ii) Special Climatological Stations. The latitudes, longitudes, and altitudes of special stations, the climatological features of which are graphically represented hereinafter, are as follows:—

Locality.	Height above Sea Level.	Latitude. S.	Longitude.	Locality.	Height above Sea Level.	Latitude.	Longitude.		
Perth Adelaide Brisbane Sydney Melbourne Hobart	Feet. 197 140 137 138 115	deg. min. 31 57 34 56 27 28 33 52 37 49 42 53	deg. min. 115 50 138 35 153 2 151 12 144 58 147 20	Darwin Daly Waters Alice Springs Dubbo Laverton, W.A. Coolgardie	Feet. 97 691 1,926 870 1,530 1,389	deg. min. 12 28 16 16 23 38 32 18 28 40 30 57	deg. min. 130 51 133 23 133 37 148 35 122 23 121 10		

SPECIAL CLIMATOLOGICAL STATIONS-AUSTRALIA.

5. Temperatures.—(i) Comparisons with other Countries. In respect of Australian temperatures generally, it may be pointed out that the isotherm for 70° Fahrenheit extends in South America and South Africa so far south as latitude 33°, while in Australia it reaches only so far south as latitude 30°, thus showing that, on the whole, Australia has latitude for latitude a more temperate climate than other places in the Southern Hemisphere.

The comparison is even more favourable when the Northern Hemisphere is included, for in the United States the 70° isotherm extends in several of the western States so far north as latitude 41°. In Europe, the same isotherm reaches almost to the southern shores of Spain, passing, however, afterwards along the northern shores of Africa till it reaches the Red Sea, when it bends northward along the eastern shore of the Mediterranean till it reaches Syria. In Asia, nearly the whole of the land area south of latitude 40° N. has a higher temperature than 70°.

The extreme range of shade temperatures in summer and winter in a very large part of Australia amounts to probably only 81°. In Siberia, in Asia, the similar range is no less than 171°, and in North America 153°, or approximately double the Australian range.

Along the northern shores of Australia the temperatures are very equable. At Darwin, for example, the difference in the means for the hottest and coldest months is only 8.3°, and the extreme readings for the year, or the highest maximum in the hottest month and the lowest reading in the coldest month, show a difference of under 50°.

Coming southward, the extreme range of temperature increases gradually on the coast, and in a more pronounced manner inland. $\mbox{ }^{\bullet}$

(ii) Hottest and Coldest Parts. A comparison of the temperatures recorded at coast and inland stations shows that, in Australia, as in other continents, the range increases with increasing distance from the coast.

In the interior of Australia, and during exceptionally dry summers, the temperature occasionally reaches or exceeds 120° in the shade, and during the dry winters the major portion of the country to the south of the tropics is subject to ground frosts. An exact knowledge of temperature disposition cannot be determined until the interior becomes more settled, but from data procurable it would appear that the hottest area of the continent is situated in the northern part of Western Australia about the Marble Bar and Nullagine goldfields, where the maximum shade temperature during the summer sometimes exceeds 100° continuously for days and weeks. The coldest part of Australia is the extreme south-east of New South Wales and extreme east of Victoria—the region of the Australian Alps. Here the temperature seldom, if ever, reaches 100°, even in the hottest of seasons.

Tasmania as a whole enjoys a most moderate and equable range of temperature throughout the year, although occasionally hot winds may cross the Straits and cause the temperature to rise to 100° in the low-lying parts.

(iii) Monthly Maximum and Minimum Temperatures. The normal monthly maximum and minimum temperatures can be best shown by means of graphs, which exhibit the

nature of the fluctuation of each for all available years. In the diagram herein for nine representative places in Australia, the upper heavy curves show the mean maximum, and the lower heavy curves the mean minimum temperatures based upon daily observations, while the other curves show the humidities.

6. Humidity.—After temperature, humidity is the most important element of climate, as regards its effect on human comfort, rainfall supply, and in connexion with engineering problems generally.

In this publication the absolute humidity has been graphically represented in the form of inches of vapour pressure (i.e. that portion of the barometric pressure due to vapour). It is this total quantity of moisture in the air which affects personal comfort, plays an important part in varying the density of the atmosphere, and in heating and refrigerating processes. The more commonly quoted value, called the relative humidity, refers to the ratio which the actual moisture contents of the air bear to the total amount possible if saturation existed at the given temperature, and is usually quoted as a percentage. The relative humidity is an important factor in all drying operations, but is much less important than the absolute humidity as affecting animal life.

The mean monthly vapour pressure has also been added to the tables of climatological data for the capital cities on pp. 72 to 77.

The normal monthly values of vapour pressure, it should be noted, combine to make the annual curve for this element which is comparable with the maximum and minimum temperature curves, but the relative humidities consisting as they do of the extremes for each month, do not show the normal annual fluctuation which would be about midway between the extremes.

The order of stations in descending values of vapour pressure is Darwin, Daly Waters, Brisbane, Sydney, Perth, Adelaide, Melbourne, Hobart and Alice Springs, while the relative humidity diminishes in the order, Sydney, Hobart, Darwin, Brisbane, Melbourne, Perth, Adelaide, Daly Waters and Alice Springs.

- 7. Evaporation.—(i) General. The rate and quantity of evaporation n any territory is influenced by the prevailing temperature, and by atmospheric humidity, pressure, and movement. In Australia, the question is of perhaps more than ordinary importance, since in its drier regions water has often to be conserved in "tanks"* and dams. The magnitude of the economic loss by evaporation will be appreciated from the tabular records herein, which show that the yearly amount varies from about 32 inches at Hobart to 94 inches at Alice Springs in the centre of the Continent. Over the inland districts of the Continent it has been calculated that evaporation equals the rainfall where the annual totals are about 36 inches, the variations above and below this quantity being inverse.
- (ii) Monthly Evaporation Curves. The curves showing the mean monthly evaporation in various parts of Australia disclose how characteristically different are the amounts for the several months in different localities. The evaporation for representative places is shown on the diagram herein.
- (iii) Loss by Evaporation. In the interior of Australia the possible evaporation is greater than the actual rainfall. Since the loss by evaporation depends largely on the exposed area, tanks and dams so designed that the surface shall be a minimum are advantageous. Further, the more protected from the direct rays of the sun and from winds, by means of suitable tree planting, the less will be the loss by evaporation. These matters are naturally of more than ordinary concern in the drier districts of Australia.
- 8. Rainfall.—(i) General. As even a casual reference to climatological maps indicating the distribution of rainfall and prevailing direction of wind would clearly show, the rainfall of any region is determined mainly by the direction and route of the prevailing winds, by the varying temperatures of the earth's surface over which they blow, and by the physiographical features generally.

Australia lies within the zones of the south-east trades and prevailing westerly winds. The southern limit of the south-east trade strikes the eastern shores at about 30° south latitude, and, with very few exceptions, the heaviest rains of the Australian continent are precipitated along the Pacific slopes to the north of that latitude, the varying quantities being more or less regulated by the differences in elevation of the shores and of the chain of mountains upon which the rain-laden winds blow from the New South Wales northern

[•] In Australia, artificial storage ponds or reservoirs are called "tanks."

border to Thursday Island. The converse effect is exemplified on the north-west coast of Western Australia, where the prevailing winds, blowing from the interior of the continent instead of from the ocean, result in the lightest coastal rain in Australia.

The westerly winds, which skirt the southern shores, are responsible for the very reliable, although generally light to moderate, rains enjoyed by the south-western portion of Western Australia, by the south-eastern agricultural areas of South Australia, by a great part of Victoria, and by the whole of Tasmania.

- (ii) Factors determining Distribution and Intensity of Rainfall. (iii) Time of Rainfall. In Official Year Book No. 6 (see pp. 72 to 74) some notes were given of the various factors governing the distribution, intensity, and period of Australian rainfall.
- (iv) Wettest and Driest Regions. The wettest known part of Australia is on the north-east coast of Queensland, between Port Douglas and Cardwell, where three stations situated on, or adjacent to, the Johnstone and Russell Rivers have an average annual rainfall of between 144 and 165 inches. The maximum and minimum falls there are:—Goondi, 241.53 in 1894 and 67.88 inches in 1915, or a range of 173.65 inches; Innisfail, 211.24 in 1894 and 69.87 inches in 1902, or a range of 141.37 inches: Harvey Creek, 254.77 in 1921 and 80.47 inches in 1902, or a range of 174.30 inches.
- On four occasions more than 200 inches have been recorded at Goondi, the last of these being in 1910, when 204.82 inches were registered. The record at this station covers a period of 32 years.

Harvey Creek, in the shorter period of 24 years has three times exceeded 200 inches, the total for 1921 being 254.77 inches, and at the South Johnstone Sugar Experiment Station, where a gauge has recently been established; 202.52 inches were recorded in 1921.

The driest known part of the continent is about the Lake Eyre district in South Australia (the only part of the continent below sea level), where the annual average is but 5 inches, and where the fall rarely exceeds 10 inches for the twelve months.

The inland districts of Western Australia were at one time regarded as the driest part of Australia, but authentic observations in recent years over the settled districts in the east of that State show that the annual average is from 10 to 12 inches.

(v) Quantities and Distribution of Rainfall. The departure from the normal rainfall increases greatly and progressively from the southern to the northern shores of the continent, and similarly also at all parts of the continent subject to capricious monsoonal rains, as the comparisons hereunder will show. The general distribution is best seen from the rainfall map herein which shows the areas subject to average annual rainfalls lying between certain limits. The areas enjoying varying quantities of rainfall determined from the latest available information are shown in the following table:—

Average Annual Rainfall.	N.S.W.	Victoria.	Queens-	South	Northern	Western Australia.	Tas.	Total.
10000000	(a)		2404.	Austrana.	remony	11 (1502 4714.	(b)	(b)
	i						i	
	sqr. mls.	sqr. mls.	sqr. mls.					
Under 10 inches	44,997	nil	91,012	317,600	138,190	513,653	nil	1,105,452
10—15 ,,	77,268	19,912	87,489	33,405	141,570	232,815	nil	592,459
15—20 ,,	57,639	12,626	112,738	14,190	62,920	89,922	937	350,972
20—30 ,,	77,202	29,317	213,779	13,827	93,470	95,404	7,559	530,558
3040 ,,	30,700	14,029	69,880	984	40,690	40,750	4,588	201,621
Over 40 ,,	22,566	12,000	95,602	64	46,780	3,376	10,101	190,489
	!						<u> </u>	
Total area	310,372	87,884	670,500	380,070	523,620	975,920	26,215	2,974,581

AVERAGE ANNUAL RAINFALL DISTRIBUTION.

Referring first to the capital cities, the complete records of which are given in the next table, it will be seen that Sydney, with a normal rainfall of 47.73 inches, occupies the chief place; Brisbane, Perth, Melbourne, Hobart and Adelaide following in that order, Adelaide with 21.20 inches being the driest. The extreme range from the wettest to the driest year is greatest at Brisbane (72.09 inches) and least at Adelaide (19.48 inches).

In order to show how the rainfall is distributed throughout the year in various parts of the continent, the figures of representative towns have been selected. (See map.)

⁽a) Including Federal Capital Territory. (b) Over an area of 3,030 square miles no records are available.

The figures for Darwin, typical of the Northern Territory, show that nearly the whole of the rainfall occurs there in the summer months, while little or none falls in the middle of the year. The figures for Perth, as representing the south-western part of the continent, are the reverse, for while the summer months are dry, the winter ones are very wet. In Melbourne and Hobart the rain is fairly well distributed throughout the twelve months, with a maximum in October for the former, and in November for the latter. The records at Alice Springs and Daly Waters indicate that in the central parts of Australia the wettest months are in the summer and autumn. In Queensland, as in the Northern Territory, the heaviest rains fall in the summer months, but good averages are also maintained during the other seasons.

On the coast of New South Wales, the first six months of the year are the wettestwith a maximum in the autumn; the averages during the last six months are fair, and moderately uniform. Generally it may be said that approximately one-third of the area of the continent, principally in the eastern and northern parts, enjoys an annual average rainfall of from 20 to 50 or more inches, the remaining two-thirds averaging from about 10 to 20 inches.

- (vi) Curves of Rainfall and Evaporation. The relative amounts of rainfall and evaporation at different times through the year are clearly indicated in the graphs herein. Inspection thereof will show how large is the evaporation when water is fully exposed to the direct rays of the sun and to wind.
- (vii) Tables of Rainfall. The table of rainfall for a long period of years for each of the various Australian capitals affords information as to the variability of the fall in successive years, and the list of the more remarkable falls furnishes information as to what may be expected on particular occasions. As pointed out in 4 ante, the capitals are dealt with in the order in which they occur in the adopted meteorological divisions.

RAINFALL-AUSTRALIAN CAPITAL CITIES, 1901 TO 1924.

]	PERT	гн.	AD	ELA	IDE.	BR	ISBA	NE.	S	ZDN1	EY.	ME	LBOU	RNE.	н	OBAI	RT.
Year.	Amount.	No. of Days.	10 Years' Means.	Amount.	No. of Days.	10 Years' Means.	Amount.	No. of Days.	10 Years' Means.	Amount.	No. of Days.	10 Years' Means.	Amount.	No. of Days.	10 Years' Means.	Amount.	No. of Days.	10 Years' Means.
		— '		<u>'</u>	_		ļ	_					i			i		
1901 2 3 4 5 6 7 8	in. 36.75 27.06 35.69 34.35 34.61 32.37 40.12 30.52 39.11	93 140 125 116 121 132 106	in.	in. 18.01 16.02 25.47 20.31 22.28 26.51 17.78 24.56 27.69	123 134 117 131 127 125 125	in.	in. 38.48 16.17 49.27 33.23 36.76 42.85 31.46 44.01 34.06	87 136 124 108 125 119 125	in.	in. 40.10 43.07 38.62 45.93 35.03 31.89 31.32 45.65 32.45	180 173 158 145 160 132 167	١	in. 27.45 23.08 28.43 29.72 25.64 22.29 22.26 17.72 25.86	102 130 128 129 114 102 130	in.	in. 25.11 21.85 25.86 22.41 32.09 23.31 25.92 16.50 27.29	150 139 139 168 155 166 148	in 23.29
1910 11 12 13 14 15 16 17 18 19 20 21 22 23 24 	37.02 23.38 27.85 38.28 20.21 43.61 35.16 45.64 39.58 30.66 40.35 41.09 31.86 44.47 33.79	108 123 141 128 164 128 146 138 120 124 135 135 134 119	34.98	24.62 15.99 19.57 18.16 11.39 19.38 28.16 28.90 17.41 17.21 26.70 22.64 23.20 29.79 23.44	127 116 102 91 117 142 153 107 108 119 100 117 139 143	21.13	49.00 35.21 41.30 40.81 33.99 25.66 52.80 40.92 24.95 19.36 39.72 54.31 35.82 23.27 41.08	128 114 115 141 93 136 127 121 96 122 167 109 93 114		46.91 50.24 47.51 57.70 56.42 34.83 44.91 52.40 42.99 58.71 43.42 43.34 39.35 37.01 37.01	155 172 141 149 117 161 151 152 159 140 123 136	46.64	24.61 36.61 20.37 21.17 18.57 20.95 38.04 30.57 27.13 24.89 28.27 29.76 25.02 22.64 36.48	168 157 157 129 167 170 171 160 141 162 154 151 158 171	26.39	25.22 26.78 23.14 19.36 15.42 20.91 43.39 30.62 26.04 22.48 18.00 18.04 28.27 32.93 28.76	193 181 165 154 196 203 214 179 153 182 159 189 198	25.82
No. of Yrs.	49	49		86	86		75	65	l 	85	85	<u></u>	81	69	_••	82	82	<u></u>

Note.—The above average rainfall figures for Brisbane, Sydney, and Melbourne differ slightly from the mean annual falls given in the Climatological Tables, which are for a less number of years. Annual totals from 1860 to 1900 inclusive will be found in Official Year Book No. 15, page 53.

9. Remarkable Falls of Rain.—The following are the more remarkable falls of rain in the various States and in the Northern Territory, which have occurred within a period of twenty-four hours. In New South Wales and Queensland falls of less than 15 inches in the twenty-four hours are not included. Reference, however, to them may be found in preceding Official Year Books (see No. 14, pp. 60-63):—

HEAVY RAINFALLS-NEW SOUTH WALES, UP TO 1924, INCLUSIVE.

Date. Am	nt. Name of Town or Locality. Date.	Amnt.
Mar., 1887 17. Feb., 1919 17. , 1898 20. Jan. 1911 20. Dec., 1898 17. , 1919 16. Feb., 1887 18. Mar. 1887 18. Feb., 1898 22.	14 Madden's Creek 13 Jan., 1911 88 Morpeth 9 Mar., 1893 13 Jan., 1911 27 Feb., 1919 14 Tongarra Farm 14 , 1898 58 South Head (near 66 Sydney) 29 Apr., 1841 58 , , 16 Oct., 1844	ins. 18.68 21.52 18.25 17.87 15.12 20.00 20.12 20.41
	Mar., 1887 Feb., 1919 ,, 1898 Jan. 1911 Dec., 1898 ,, 1919 Feb., , Mar. 1887 Feb., 1898 22.5	Mar., 1887 17.14 Madden's Creek 13 Jan., 1911 Feb., 1919 17.88 Morpeth 9 Mar., 1893 Jan. 1911 20.83 Numbugga 27 Feb., 1919 Tongarra Farm 14 , 1898 Feb., 18.58 South Head (near Mar. 1887 18.66 Sydney) 29 Apr., 1841 Feb., 1898 22.58 , , , , 16 Oct., 1844

HEAVY RAINFALLS-QUEENSLAND, UP TO 1924, INCLUSIVE.

Name of Town or Locality.	Date.	Amnt.	Name of Town or Locality.	Date.	Amnt.
Anglesey	26 Dec., 1909	ins. 18.20	Flying Fish Point	31 Jan., 1913	ins. 16.10
Atherton (Cairns)	31 Jan., 1913	16.69	Gladstone	4 Feb., 1911	18.83
Babinda (Cairns)	1 Feb., ,,	20.51	Glen Boughton	5 Apr., 1894	18.50
,, ,,	24 Jan., 1916	22.30	Goldsborough		!
,, ,,	21 Apr., 1920	16.05	(Cairns)	31 Jan., 1913	19.92
Babinda	25 Mar., 1921	15.76	Goondi Mill (Innis-	1	
Bloomsbury	14 Feb., 1893	17.40	fail)	6 Apr., 1894	15.69
,,	10 Jan., 1901	16.62	,, ,,	29 Dec., 1903	17.83
Brisbane	21 ,, 1887	18.31	,, ,,	10 Feb., 1911	17.68
Buderim Mountain	11 ,, 1898	26.20	,, ,,	6 Apr., 1912	15.55
Bundaberg	16 ,, 1913	16.94	Goondi	30 Jan., 1913	24.10
Burnett Head			Goorganga	23 , 1918	18.17
(Bundaberg)	16 , 1913	15.22	Halifax	5 Feb., 1899	15.37
Cairns	11 Feb., 1911	15.17		6 Jan., 1901	15.68
,,,	2 Apr., ,,	20.16	Hambledon Mill	2 ,, 1911	18.61
Carbrook	23 Jan., 1918	22.66	,, ,,	1 Apr., ,,	19.62
	24 ,, ,,	15.77		30 Jan., 1913	17.32
Cardwell	18 Mar., 1904	18.24	Hampden	23 Apr., 1918	17.30
Carmilla	23 Jan., 1918	15.92	,,	24 ,, ,,	17.19
Clare	26 , 1896	15.30	Harvey Creek	8 Mar., 1899	17.72
Collaroy	23 , 1918	18.06	,, ,,	11 Jan., 1905	16.96
Crohamhurst	0.77.1		,, ,,	3 ,, 1911	27.75
(Blackall Range)	2 Feb., 1893	35.71	,, ,,	2 Apr., ,,	16.46
,, ,,	9 Jan., 1898	19.55	,, ,,	31 Jan., 1913	24.72
a , " "	6 Mar., ,,	16.01	,, ,,	25 Mar., 1921	15.80
Croydon	29 Jan., 1908	15.00	,, ,,	12 Mar., 1924	16.50
Dungeness	16 Mar., 1893	22.17	,, ,,	13 Mar., ,,	15.78
Dunira	9 Jan., 1898	18.45	Haughton Valley	26 Jan., 1896	18.10
T	6 Mar., "	15.95	Holmwood (Wood-	0 T 1 1000	10.10
Fairymead Planta	10 T 1010	1, 00	ford)	2 Feb., 1893	16.19
tion (Bundaberg)	16 Jan., 1913	15.32	Howard	15 Jan., 1905	19.55
Flying Fish Point	7 Apr., 1912	16.06	Huntley	27 Dec., 1916	18.94

HEAVY RAINFALLS—QUEENSLAND—continued.

Name of Town or Locality.	Date.	Amnt.	Name of Town or Locality.	Date.	Amnt.
		ins.			ins.
Innisfail (formerly)		Mourilyan	11 Feb., 1911	17.40
Geraldton)	11 Feb., 1889	17.13	,,	7 Apr., 1912	18.97
,, ,,	6 Apr.,1894	16.02	,,	31 Jan., 1913	15.05
,, ,,	24 Jan., 1900	15.22	Mundoolun	21 ,, 1887	17.95
,, ,,	29 Dec., 1903	21.22	Nambour	9 ,, 1898	21.00
,, ,,	2 Apr., 1911	15.00	,,	27 Dec., 1909	16.80
,, ,,	7 , 1912	20.50	Netherdale	22 Jan., 1918	19.50
,, ,,	31 Jan., 1913	20.91	Oxenford	14 Mar., 1908	15.65
Kamerunga (Cairns)	2 Apr., 1911	21.00	Palmwoods	10 Jan., 1898	15.85
,	31 Jan., 1913	16.00		25 Dec., 1909	17.75
Koumala	23 , 1918	22.31	Pialba (Marybor'gh)	16 Jan., 1913	17.22
	94	20.65	Plane Creek		
Kuranda (Cairns)	11 Feb., 1911	16.30	(Mackay)	26 Feb., ,,	27.73
• •	17.35	15.10	Port Douglas	10 Mar., 1904	16.34
	01	18.60		17 , 1911	16.10
,, ,,		24.30	,, ,,	"	31.53
,, ,,	n -	28.80	Proserpine	1 Apr., ,, 23 Jan., 1918	18.17
,, ,,	31 Jan., 1913	16.34	70	24 Mar., 1890	17.00
Landsborough		15.15	70 1 11/7		17.35
T T1 1	2 Feb., 1893	15.15	D 1.	16 Feb., 1893	18.90
	10 Mar., 1904	1 1	~ .	16 Jan., 1913 23 1918	22.60
Lyndon (via Brixton)	1 Apr., 1911	15.30	Sarina	"	
Lyndon (via Brixton)	3 , 1917	17.00*	St. Lawrence	30 ,, 1896	15.00
Mackay	21 Jan., 1918	24.70†	The Hollow (Mac-	20 77 1 7000	17.10
~ " =	22 ,, ,,	17.25‡	kay)	23 Feb., 1888	15.12
Sugar Experimental	i		Thornborough	20 Apr., 1903	18.07
Farm, Mackay	21 ,, ,,	16.80	Townsville	24 Jan., 1892	19.20
,,	22 ,, ,,	17.20	,,	28 Dec., 1903	15.00
Macnade Mill	5 Feb., 1899	15.20	Victoria Mill	6 Jan., 1901	16.67
,,	6 Jan., 1901	23.33	Woodlands (Yepp'n)	31 ,, 1893	23.07
,,	4 Mar., 1915	22.00	Wootha	10 Feb., 1915	15.93
Mapleton	26 Dec., 1909	15.72	Yandina	1 ,, 1893	20.08
Mirani	12 Jan., 1901	16.59	l ",	9 Jan., 1898	19.25
Miriam Vale (B'berg)	17 ,, 1913	15.80	,,	28 Dec., 1909	15.80
Mooloolah	13 Mar., 1892	21.53	Yarrabah	2 Apr., 1911	30.65
,,	2 Feb., 1893	19.11	,,	24 Jan., 1916	27.20
Mount Cuthbert	8 Jan., 1911	18.00	,,	25 ,, ,,	18.60
Mount Molloy	31 Mar., ,,	20.00	Yeppoon	31 ,, 1893	20.05
,,	1 Apr., ,,	20.00	,,	8 ,, 1898	18.05
,,	2 ,, ,,	20.00	,,	. 8 Oct., 1914	21.70
,,	""		"		

HEAVY RAINFALLS-WESTERN AUSTRALIA, UP TO 1924, INCLUSIVE.

Name of Town Locality.	or	Date.	Amnt.	Name of Town of Locality.	or	Date.	Amnt.
Alice Downs Balla Balla Bamboo Creek Boodarie Booloogooroo Broome Carlton Cossack Croydon Derby ,		15 Mar., 1922 21 ,, 1899 22 ,, ,, 3 Jan., 1896 16 ,, 1923 6 Jan., 1917 11 ,, 1903 3 Apr., 1898 16 ,, 1900 3 Mar., 1903 29 Dec., 1898 7 Jan., 1917	ins. 10.58 14.40 10.10 14.53 10.03 10.76 14.00 10.64 12.82 13.23 12.00 13.09 16.47	Exmouth Gulf Fortescue Frazier Downs Gnaraloo Kerdiadary Meda Millstream Minilya Obagama "Pilbara Point Cloates		2 Feb., 1918 3 May, 1890 3 Mar., 1916 20 , 1923 7 Feb., 1901 2 Mar., 1916 5 ,, 1900 15 Jan., 1923 28 Feb., 1910 24 Dec., 1920 2 Apr., 1898 20 Jan., 1909	10.8. 12.50 23.36 11.25 11.00 12.00 10.55 10.00 11.50 12.00 13.02 14.04 10.87

[•] Mr. Jas. Laidlaw, of Lyndon, states that this fell in 4 hours. † 371 hours. ‡ 221 hours.

HEAVY RAINFALLS, WESTERN AUSTRALIA -continued.

Pgint Torment	Name of Town or Locality.		Date.	Amnt.	Name of Town Locality.	or	Date.	Amnt.
Whim Creek 3 Apr., 1898 29.41	Port George IV. Roebourne Roebuck Plains Springvale Tambray Thangoc		17 Jan., 1915 3 Apr., 1898 6 Mar., 1900 5 Jan., 1917 6 ", 1900 3 ", 1903 17-19 Feb. '96 28 Dec., 1898	11.86 11.24 11.44 10.32 14.01 22.36 12.25 11.00 10.47 24.18 11.15	Winderrie Woodstock Wyndham Yardie Creek Yeeda		6 ,, 1900 3 ,, 1903 17 Jan., 1923 21 ,, 1912 27 Jan., 1890 4 Mar., 1919 3 Feb., 1918 2 Mar., 1916 6 Jan., 1917	18.17 10.03 10.44 14.23 13.00 11.60 12.50 10.00 10.70 10.20

HEAVY RAINFALLS-NORTHERN TERRITORY, UP TO 1924, INCLUSIVE.

	Name of Town or Locality.		Amnt.	Name of Town or Locality.	Date.	Amnt.
Bonrook Borroloola Brock's Creek	 Gold	24 Dec., 1915 14 Mar., 1899 4 Jan., 1914 24 Dec., 1915 4 Jan., 1914 24 Dec., 1915	ins. 10.60 14.00 10.68 14.33 11.61	Darwin	7 Dec., 1915 30-31 Mar.,'23 21 Mar., 1901 8 Jan., 1897	ins. 11.67 12.00a 10.25 10.35

⁽a) Approximate only, as gauge was washed away.

HEAVY RAINFALLS—SOUTH AUSTRALIA, UP TO 1924, INCLUSIVE.

Name of Town or Locality.	Date.	Amnt.	Name of Town or Locality.	Date.	Amnt.
Wilmington	28 Feb., 1921	ins. 3.97	Wilmington	1 Mar., 1921	ins. 7.12

HEAVY RAINFALLS-VICTORIA, UP TO 1924, INCLUSIVE.

	of Tow ocality.		Dat	e. 	Amnt.	Name of Tov Locality		Date	•	Amnt.
Balook		 26 8 27 28	Sept.	,1917 ,,	ins. 5.32 7.23 2.08	Mt. Buffalo	••	6 June, 7 ,,	1917	ins. 8.53 6.56

HEAVY RAINFALLS—TASMANIA, UP TO 1924, INCLUSIVE.

Name of Town or Locality.	Date.	Amnt.	Name of Town or Locality.	Date.	Amnt.
Cullenswood Gould's Country Lottah	5 June, 1923 8-10 Mar., '11 8-10 ,, ,,	15.33	The Springs	8-10 Mar., '11 30-31 Jan., '16 5 June, 1923	10.75

10. Snowfall.—Light snow has been known to fall occasionally so far north as latitude 31° S., and from the western to the eastern shores of the continent. During exceptional seasons, it has fallen simultaneously over two-thirds of the State of New South Wales, and has extended at times along the whole of the Great Dividing Range, from its southern extremity in Victoria so far north as Toowoomba in Queensland. During the winter, for several months, snow covers the ground to a great extent on the Australian Alps, where also the temperature falls below zero Fahrenheit during the night. In the ravines around Kosciusko and similar localities the snow never entirely disappears.

The antarctic "V"-shaped disturbances are always associated with the most pronounced and extensive snowfalls. The barometric gradients are very steep where the "trough line" extends northward, and the apexes are unusually sharp-pointed, and protrude into very low latitudes, sometimes even to the tropics.

11. Hail.—Hail falls most frequently along the southern shores of the continent in the winter, and over south-eastern Australia during the summer months. The size of the hailstones generally increases with distance from the coast, a fact which lends strong support to the theory that hail is brought about by ascending currents. A summer rarely passes without some station experiencing a fall of stones exceeding in size an ordinary hen-egg, and many riddled sheets of light-gauge galvanized iron bear evidence of the weight and penetrating power of the stones.

The hailstorms occur most frequently when the barometric readings indicate a flat and unstable condition of pressure. They are almost invariably associated with tornadoes or tornadic tendencies, and on the east coast the clouds from which the stones fall are generally of a remarkable sepia-coloured tint.

- 12. Barometric Pressures.—The mean annual barometric pressure (corrected to sealevel and standard gravity) in Australia varies from 29.80 inches on the north coast to 29.92 inches over the central and 30.03 inches in the southern parts of the continent. In January, the mean pressure ranges from 29.70 inches in the northern and central areas to 29.95 inches in the southern. The July mean pressure ranges from 29.90 inches at Darwin to 30.12 inches at Alice Springs. Barometer readings corrected to mean sea level and standard gravity have, under anticyclonic conditions in the interior of the continent, ranged as high as 30.77 inches (at Kalgoorlie on the 28th July, 1901) and have fallen as low as 27.55 inches. This lowest record was registered at Mackay during a tropical hurricane on the 21st January, 1918. An almost equally abnormal reading of 27.88 inches was recorded at Innisfail during a similar storm on the 10th March, 1918. The mean annual fluctuations of barometric pressure for the capitals of Australia are shown on the graph herein.
- 13. Wind.—Notes on the distinctive wind currents in Australia were given in preceding Year Books (see No. 6, page 83), but, owing to limitations of space, have not been included herein.
- 14. Cyclones and Storms.—The "elements" in Australia are ordinarily peaceful, and while destructive cyclones have visited various parts, more especially coastal areas, such visitations are rare, and may be properly described as erratic.

During the winter months, the southern shores of the continent are subject to cyclonic storms, evolved from the V-shaped depressions of the southern low-pressure belt. They are felt most severely over the south-western parts of Western Australia, to the southeast of South Australia, in Bass Straits, including the coast line of Victoria, and on the west coast of Tasmania. Apparently the more violent wind pressures from these cyclones are experienced in their northern half, or in that part of them which has a north-westerly to a south-westerly circulation.

The north-east coast of Queensland is occasionally visited by hurricanes from the north-east tropics. During the first four months of the year, these hurricanes appear to have their origin in the neighbourhood of the South Pacific Islands, their path being a parabolic curve first to the S.W. and finally towards the S.E. Only a small percentage, however, reach Australia, the majority recurving in their path to the east of New Caledonia.

Very severe cyclones, locally known as "willy willies," are peculiar to the northwest coast of Western Australia from the months of November to April inclusive. They apparently originate in the ocean in the vicinity of Cambridge Gulf, and travel in a southwesterly direction with continually increasing force, displaying their greatest energy near Cossack and Onslow, between latitudes 20° and 22° South. The winds in these

storms, like those from the north-east tropics, are very violent and destructive, and cause great havoc amongst the pearl-fishers. The greatest velocities are usually to be found in the south-eastern quadrant of the cyclones, with north-east to east winds. After leaving the north-west coast, these storms either travel southwards, following the coastline, or cross the continent to the Great Australian Bight. When they take the latter course, their track is marked by torrential rains, as much as 29.41 inches, for example, being recorded in 24 hours at Whim Creek from one such occurrence. Falls of 10 inches and over have frequently been recorded in the northern interior of Western Australia from similar storms.

Some further notes on severe cyclones and on "southerly bursters," a characteristic feature of the eastern part of Australia, will be found in previous issues of the Official Year Book (see No. 6, pp. 84, 85, 86).

A special article dealing with "Australian Hurricanes and Related Storms" appeared in Official Year Book No 16, pp. 80-84.


- 15. Influences affecting Australian Climate.—(i) General. Australian history does not cover a sufficient period, nor is the country sufficiently occupied, to ascertain whether or not the advance of settlement has materially affected the climate as a whole. Local changes have, however, taken place, a fact which suggests that settlement and the treatment of the land have a distinct effect on local conditions. For example, the mean temperature of Sydney shows a rise of two-tenths of a degree during the last twenty years, a change probably brought about by the great increase of residential and manufacturing buildings within the city and in the surrounding suburbs. Again, low-lying lands on the north coast of New South Wales, which originally were seldom subject to frosts, have, with the denudation of the surrounding hills from forests, experienced annual visitations, the probable explanation being that through the absence of trees the cold air of the high lands now flows unchecked and untempered down the sides of the hills to the valleys and lower lands.
- (ii) Influence of Forests on Climate. As already indicated, forests doubtless exercise a great influence on local climate, and hence, to the extent that forestal undertakings will allow, the weather can be controlled by human agency. The direct action of forests is an equalizing one; thus, especially in equatorial regions, and during the warmest portion of the year, they considerably reduce the mean temperature of the air. They also reduce the diurnal extremes of shade temperatures by altering the extent of radiating surface, by evaporation and by checking the movement of air, and while decreasing evaporation from the ground, they increase the relative humidity. Vegetation greatly diminishes the rate of flow-off of rain and the washing away of surface soil, and when a region is protected by trees, a steadier water supply is ensured, and the rainfall is better conserved. In regions of snowfall, the supply of water to rivers is similarly regulated, and without this and the sheltering influence of ravines and "gullies," watercourses supplied mainly by melting snow would be subject to alternate periods of flooding and dryness. This is borne out in the case of the inland rivers; the River Murray, for example, which has never been known to become dry, deriving its steadiness of flow mainly through the causes indicated.
- (iii) Direct Influence of Forests on Rainfall. Whether forests have a direct influence on rainfall is a debatable question, some authorities alleging that precipitation is undoubtedly induced by forests, while others take the opposite view.

Sufficient evidence exists, however, to prove that, even if the rainfall has not increased, the beneficial climatic effect of forest lands more than warrants their protection and extension. Rapid rate of evaporation, induced by both hot and cold winds, injures crops and makes life uncomfortable on the plains, and, while it may be doubted that the forest aids in increasing precipitation, it must be admitted that it does check winds and the rapid evaporation due to them. Trees as wind-breaks have been successfully planted in central parts of the United States, and there is no reason why similar experiments should not be successful in many parts of the treeless interior of Australia. The belts should be planted at right angles to the direction of the prevailing parching winds, and if not more than half a mile apart will afford shelter to the enclosed areas.

In previous issues some notes on observations made in other countries were added (see Official Year Book No. 6, pp. 86 and 95).

16. Rainfall and Temperature, Various Cities.—The following table shows rainfall and temperature for various important cities throughout the world, for the site of the Federal capital, and for the capitals of the Australian States.

ANNUAL FLUCTUATIONS OF NORMAL MAXIMUM AND MINIMUM TEMPERATURE AND HUMIDITY.



EXPLANATION.—The upper and lower heavy lines in each graph represent the maximum and minimum temperatures respectively. The Fahrenheit temperature scales are shown on the outer edge of the sheet under ",F," and the centigrade scales in the two inner columns under "C."

The broken line shows the normal absolute humidity in the form of 9 a.m. vapour pressures for which the figures in the outer "F" columns represent hundredths of an inch of barometric pressure.


The upper and lower fine lines join the greatest and the least monthly means of relative humidity respectively, the figures under the outer columns "F" indicating percentage values.


The curves for temperature and vapour pressure joining the mean monthly values serve to show the annual fluctuation of these elements, but the relative humidity graphs joining the extreme values for each month do not indicate any normal annual variation.

Comparison of the maximum and minimum temperature curves affords a measure of the mean diurnal range of temperature. At Perth in the middle of January, for instance, there is normally a range of 21° from 63° F. to 84° F., but in June it is only 15° from 48° F. to 63° F.


The relative humidity curves illustrate the extreme range of the mean monthly humidity over a number of years.

MEAN MONTHLY RAINFALL AND EVAPORATION.






Area affected and period of duration of the Longest Heat Waves when the Maximum Temperature for consecutive 24 hours reached or exceeded 100° Fah.



Greatest number of consecutive days on which the Shade Temperature was over 100° Fah. at the places indicated.



EXPLANATION.—On the preceding graphs thick lines denote rainfall, and thin lines evaporation, and show the fluctuation of the mean rate of fall *per month* throughout the year. The results, plotted from the Climatological Tables herein are shown in inches (see the outer columns), and the corresponding metric scale (centimetres) is shown in the two inner columns. The evaporation is not given for Darwin and Daly Waters

At Perth, Adelaide, Brisbane, Melbourne, Hobart, Alice Springs, and Coolgardie the results have been obtained from jacketed tanks sunk in the ground. At Sydney and Dubbo sunken tanks without water jackets are used, whilst at Laverton (W.A.) the records are taken from a small portable jacketed evaporation dish of 8 inches in diameter.


The distance for any date from the zero line to the curve represents the average number of inches, reckoned as per month, of rainfall at that date. Thus, taking the curves for Adelaide, in the middle of January the rain falls on the average at the rate of about three-fourths of an inch per month, or, say, at the rate of about 9 inches per year. In the middle of June it falls at the rate of a little over 3 inches per month, or, say, at the rate of about 37 inches per year. At Dubbo the evaporation is at the rate of nearly 11½ inches per month about the middle of January, and only about 1½ inches at the middle of June

The mean annual rainfall and evaporation at the places indicated are given in the appended table.

Evapora-Evapora-Place. Rainfall. Place. Rainfall. tion. tion. In. In. In. In. 62.07 26.58 11.24 21.82 Perth 34.10 21.18 65.80 Darwin Daly Waters ... Alice Springs ... Dubbo ... Adelaide 54.55 . . 45.20 47.74 53.37 38.43Brisbane ٠. 93.99Sydney 66.37 25.61 Melbourne 38.90 Laverton, W.A. 9.76 141.45 . . Hobart Coolgardie 10.07 87.69 .. }


MEAN ANNUAL BAINFALL AND EVAPORATION.


MEAN BAROMETRIC PRESSURE-CAPITAL CITIES.



EXPLANATION.—The lines representing the yearly fluctuations of barometric pressure at the State capital cities are means for long periods, and are plotted from the Climatological Tables herein. The pressures are shown in inches on about 2½ times the natural scale, and the corresponding pressures in centimetres are also shown in the two inner columns, in which each division represents one millimetre.

INTERPRETATION.—Taking the Brisbane graph for purposes of illustration, it will be seen that the mean pressure in the middle of January is about 29.87 inches, and there are maxima in the middle of May and August of about 30.09 inches.





RAINFALL AND TEMPERATURE-VARIOUS CITIES.

•		An	nual Rain	fall.			Tempe	rature.		
Place.	Height above M.S.L.	Average.	Highest.	Lowest.	(a)Mean Summer.	(b)Mean Winter.	Highest on Record.	Lowest on Record.	Average Hottest Month.	Average Coldest Month.
	Ft.	Ins.	Ins.	Ins.	Fahr.	Fahr.	Fahr.	Fahr.	Fahr.	Fahr.
Amsterdam	6 125	27.29	40.59	17.60 26.32	63.2	36.8	90.0	4.1	64.4	35.4
Auckland	351	43.88 15.48	74.15 33.33	4.56	66.2 79.2	52.5 49.1	91.0 109.4	31.9 19.6	67.1 81.0	51.8 47.4
Rergen	72	77 00	111.58	44.49	56.8	34.2	88.5	4.8	57.9	33.6
Berlin	161	77.09 22.72 36.30	30.04	14.25	64.8	34.2 33.0	98.6	-13.0	66.0	31.8
Berne	1,877	36.30	30.04 58.23	24.69	162.2	30.1	91.4	-3.6	64.4	28.0
Bombay	37	71.15	114.89	33.41	83.5	75.1	100.0	55.9	84.8	74.2
Authens	482 328	22.52 28.35	32.56 41.18	16.50	64.1	33.5 36.0	100.0 95.5	$\begin{bmatrix} -23.4 \\ -4.4 \end{bmatrix}$	65.5 63.7	29.3 34.5
Dadonest	500	25.20	35.28	17.73 16.79	68.6	30.2	98.6	- 5.1	70.4	28.2
Buenos Ayres Calcutta	82	25.20 38.78	35.28 79.72 98.48	20.04	72.7	50.9	103.1	$^{-5.1}_{22.3}$	73.8	50.0
Calcutta	21	61.82	98.48	38.43	85.6	68.0	108.2	44.2	86.0	66.4
Capcionii	40	25.50	36.72	17.71	68.1	54.7 65.3	102.0	34.0	68.8 69.2	53.9
Caracas Chicago	3,420 823	30.03	47.36 45.86	23.70 24.52	68.3	26.1	87.8 103.0	$^{48.2}_{-23.0}$	79.2	$63.7 \\ 23.7$
Christehurch	25	33.28 25.16	35.30	13.54	70.0 61.3	43.3	95.7	21.3	72.4 61.6	42.4
Christiania (Oslo)	75	23.23	35.30 32.21	13.54 16.26	61.0	24.5	95.0	-21.1	62.6	23.9
Colombo	40	83.83	139.70	51.60	81.5	79.9	95.8	65.0	82.6	79.1
Constantinople	245	28.75	42.74	14.78	74.0	43.5	103.6	13.0	75.7	42.0
Copenhagen Dresden	10 115	20.79	25.83 34.49	15.47 17.72 16.60	62.9	33.3 32.4	85.5 93.4	$-3.3 \\ -15.3$	61.9	32.4 31.6
Dark Ray	47	26.80 27.66	35.56	16.60	59.4	42.0	87.2	13.3	60.5	41.7
Dunedin	300	36.96	54.51	22.15	56.3	42.6	94.0	23.0	57.0	41.5
Durnan	260	40.79	71.27	27.24	75.6	64.4	110.6	41.1	76.7	63.8
Edinburgh Geneva	441	25.21	32.05	16.44	55.8	38.8	87.7	5.0	57.2	38.3 32.2
Genoa	1,328 157	33.48 51.29	46.89 108.22	21.14 28.21	64.4 73.8	33.7 46.8	94.5	16.7	66.2 75.4	45.5
Genoa	184	38.49	56.18	29.05	52.7	41.0	84.9	6.6	58.0	38.4
Greenwich	149	23.50	35.54 119.72	16.38	62.0	39.5	100.0	6.9	63.5	38.5
Hong Kong	109	84.28	119.72	45.84	86.2	64.8	97.0	32.0 23.3	63.5 86.7 68.2	62.9
Johannesburg	5,750	31.63	50.00	21.66	65.4	54.4 31.5	94.0	-14.8	64.8	48.9 30.6
Leipzig	384 312	24.69 29.18	$ \begin{array}{r} 31.37 \\ 52.79 \end{array} $	21.66 17.10 17.32	63.1 69.6	51.3	$97.3 \\ 94.1$	32.5	70.2	49.3
	18	23.80	38.20	16.64	61.2	39.8	94.0	9.4	62.7	38.9
Madras	22	49.85	88.41	18.45	89.0	76.8	113.0	57.5	89.9	76.1
Madrid	2,149	16.23	27.48	$9.13 \\ 12.28$	73.0	41.2	107.1	10.5 11.7	75.7 72.3	39.7 44.6
Moscow	246 526	22.24 18.94	43.03 29.28	12.28	70.5 63.4	45.3 14.7	$\frac{100.4}{99.5}$	-44.5	66.1	11.9
Naples	489	34.00	56.58	21.75	73.6	48.0	99.1	23.9	75.4	46.8
New York	314	44.63	56.58 58.68	33.17	71.4	31.8	102.0	-13.0	73.5	30.2
Ottawa	236	33.40	53.79	25.63	67.2	14.1	98.0	-33.0	69.7	12.0
Paris	164	22.64	29.57 36.00	16.46	63.5 77.7	37.2 26.6	$101.1 \\ 114.0$	-14.1 - 5.0	64.9 79.2	36.1 23.6
Petrograd	143 16	$\frac{24.40}{21.30}$	29.52	18.00 13.75	61.1	17.4	97.0	-38.2	63.7	15.2
Pekin Petrograd Quebec Rome San Francisco	16 296	40.50	53.79	32.12	63.5	12.4	96.0	-34.0	66.3	10.1
Rome	166	32.57	57.89	12.72	74.3	46.0	104.2	17.2	76.1	44.6
San Francisco	155	22.27	38.82 62.52	9.00 27.92	58.8	50.5	101.0 102.9	29.0	59.3 80.4	49.5 37.8
Shanghai	21 8	45.00 91.99	158.68	32.71	78.0 81.2	41.1 78.6	94.2	10.2 63.4	81.5	78.3
Shanghai Singapore Stockholm	144	19.09	28.27	11.81	59.5	27.3	96.8	-25.6	61.9	26.4
	65	61.45	86.37	45.72 26.57	74.8	39.2	97.9	17.2	77.7	37.5
Trieste	85	$\frac{42.94}{24.50}$	63.14 33.90	26.57	73.9	41.3	99.5 97.7	14.0	76.3	39.9
Vienna Vladivostock	663 55	$\frac{24.50}{19.54}$	33.90 33.60	$\frac{16.50}{9.39}$	65.7 63.9	30.4 11.0	97.7	-8.0 -21.8	67.1 69.4	28.0 6.1
Vladivostock Washington	112	43.50	61.33	30.85	74.7	34.5	106.0	-15.0	76.8	32.9
Wellington (N.Z.)	10 l	48.65	67.68	27.83	61.8	48.6	88.0	28.6	62.5	47.7
Zürich	1,542	45.15	78.27	29.02	63.3	31.3	94.1	1 - 0.8	65.1	29.5
]	FEDERAI	CAPIT	AL SIT					
Canberra (Dist.) Queanbeyan	${ 2,000 \atop to \atop 2,900} $	22.50	41.29	10.45	(a) 68.0	(b) 44.0	102.6	14.0	68.6	42.7
			STAT	e Capi	TALS.					
					(a)	(b)		1	1	1
	197	34.08	46.73	20.21	73.0	56.0	108.4	34.2	74.2	55.2
Perth			000-							
Perth Adelaide	140	21.21	30.87	11.39	72.6	53.1	116.3	32.0	74.1	51.7
Perth	140 137	21.21	30.87	16.17 21.49	76.7	59.7	108.9 108.5	36.1 35.9	74.1 77.2 71.7	58.5 52.6
Perth	140		30.87 88.26 82.76 44.25 43.39	11.39 16.17 21.49 15.61 13.43	72.6 76.7 71.0 66.6 61.6	59.7 54.1 50.0	108.9 108.5 111.2	36.1 35.9 27.0 27.0	74.1 77.2 71.7 67.5 62.3	51.7 58.5 52.6 48.6 45.6

⁽a) Mean of the three hottest months.

17. Climatological Tables.—The means, averages, extremes, totals, etc., for a number of climatological elements have been determined from long series of observations at the Australian capitals up to and including the year 1924. These are given in the following tables:—

⁽b) Mean of the three coldest months.

CLIMATOLOGICAL DATA-PERTH, WESTERN AUSTRALIA.

Lat. 31° 57′ S., Long. 115° 50′ E. Height above M.S.L. 197 Ft.

BAROMETER, WIND, EVAPORATION, LIGHTNING, CLOUDS, AND CLEAR DAYS.

	ed Sea tan- y and ngs.		Wine	d.		# g		ount 9 a.m.	[
Month.	Bar. corrected to 32° F. Mn. Sec Level and Stan- dard Gravity from 9 a.m. and 3 p.m. readings.	Greatest Number of Miles in one day.	Mean Hourly Pres- sure. (lbs.)	Total Miles.	Prevailing- Direction.	Mean Amount of Evaporation (inches).	No. of Days Lightning.	Mean Amount of Clouds, 9 a.1 3 p.m. & 9 p.n	No. of Clear Days.
No. of yrs. over which observation extends	40	27	27	27	27	26	27	28	28
January February March	29.908 29.924 29.987	797 27/98 650 6/08 651 6/13	0.67 0.63 0.53	11,192 9,786 9,950	SSE SSE SSE	10.35 8.59 7.65	1.7 1.5 1.4	2.9 2.9 3.2	13.9 11.6 12.0
April May	30.075 30.073 30.058	955 25/00 768 5/12 861 27/10	0.39 0.34 0.35	8,343 8,073 7,966	SE ENE N	4.75 2.72 1.75	1.3 2.3 2.2	4.2 5.3 5.9	7.9 5.1 3.2
July August September	30.092 30.086 30.060	949 11/99 966 15/03 864 11/05	0.38 0.40 0.42	8,485 8,803 8,618	N W S W	1.75 2.35 3.30	2.4 1.6 1.3	5.4 5.4 4.9	5.0 5.0 5.7
October November	30.030 29.988 29.927	809 6/16 777 18/97 776 6/22	0.52 0.58 0.63	9,876 10,099 10,955	ssw s	5.22 7.60 9.77	1.0 1.3 1.8	4.9 3.8 2.9	5.9 8.1 12.3
Year { Totals Averages Extremes	30.017	966 15/8/03	0.48	9,346		65.80	19.8	4.3	95.7

TEMPERATURE AND SUNSHINE.

		n Tem re (Fal			e Shade ire (Fahr.).	e .	Extr Temperat	eme ure (Fahr.).	ine.
Month.	Mean Max	Mean Min.	Mean.	Highest.	Lowest.	Extreme Range.	Highest in Sun.	Lowest on Grass.	Mean Hours of Sunshine.
No. of yrs. over whice observation extend		28	28	28	28	28	26	26	27
February	. 84.3 . 85.0 . 81.5 . 76.2	63.1 63.5 60.9 57.2	73.7 74.2 71.2 66.7	108.4 28/21 107.4 4/23 106.4 14/22 99.7 9/10	49.9 1/21 47.7 1/02 45.8 8/03 39.3 20/14	58.5 59.7 60.6 60.4	177.3 22/14 169.0 4/99 167.0 19/18 157.0 8/16	40.4 1/21 39.8 1/13 36.7 8/03 31.0 20/14	318.4 270.8 269.5 218.7
June July August	68.7 63.9 62.7 63.8	52.5 49.5 47.8 48.1	60.6 56.7 55.2 56.0	90.4 2/07 81.7 2/14 76.4 21/21 81.0 12/14	34.3 11/14 35.0 30/20 34.2 7/16 35.3 31/08	56.1 46.7 42.2 45.7	144.2 8/24 135.5 9/14 133.2 13/15 145.1 29/21	25.3 11/14 26.5 30/20 25.1 30/20 27.9 10/11	176.5 144.2 166.7 184.9
October	. 66.0 . 69.1 . 75.5 . 80.7	50.1 52.6 56.5 60.4	58.0 60.8 66.0 70.6	90.9 30/18 95.3 30/22 104.6 24/13 107.9 20/04	38.9 17/13 40.5 5/24 42.0 1/04 48.0 2/10	52.0 54.8 62.6 59.9	153.6 29/16 154.0 29/14 166.6 23/15 168.7 25/15	29.2 21/16 30.5 4/17 35.5 (a) 39.0 12/20	203.2 235.9 285.7 324.4
I car Durtnomen	73.1	55.2	64.1	 108.4 28/1/21	34.2 7/7/16	74.2	177.3 22/1/14	25.1 30/7/20	2798.95

(a) 6/10 and 14/12.

(b) Total for year.

	Vapour Pressure	Rel.	Hum.	(%)		Rainfall (inches).							
Month.	Mean 9 a.m.	Mean 9 a.m.	Highest Mean.	Lowest Mean.	Mean Monthly.	Mean No. of Days Rain.	Greatest	Monthly.	Least	Monthly.		in One Day.	Mean No. Days Dew.
No. of yrs. over which observation extends	23	28	28	28	49	49	4	9	.4	19		49	28
January February March April May June July August September October November December	.440 .443 .423 .398 .368 .339 .318 .325 .3 0 .350 .380 .417	52 54 56 63 73 78 77 74 68 62 55	61 65 66 72 81 83 84 79 75 63 62	42 46 46 53 61 68 69 67 58 54 46	0.34 0.45 0.76 1.59 4.94 6.89 6.44 5.70 3.40 2.18 0.80 0.59	3 2 4 7 14 17 17 18 15 12 6 4	2.17 2.98 4.50 4.97 12.13 12.80 11.29 10.33 7.84 7.87 2.78 3.05	1879 1915 1896 1882 1879 1922 1917 1882 1922 1890 1916 1888	nil nil nil nil 0.98 2.16 2.42 0.46 0.34 0.45 nil	(a) (a) (a) 1920 1903 1877 1876 1902 1916 1892 1891 1886	1.74 1.63 2.06 2.62 2.80 3.90 3.00 2.79 1.73 1.38 1.11	28/79 26/15 26/23 30/04 20/79 10/20 4/91 7/03 23/09 15/10 30/03 1/88	2.6 3.2 5.6 9.1 12.4 12.2 13.3 11.3 9.3 5.3 3.8 3.0
Year { Totals Averages Extremes	.375	62	 84		34.08	119	12.80 6/22				3.90 10/6/2		91.1

⁽a) Various years.

⁽b) January, February, March, November, and December, various years.

CLIMATOLOGICAL DATA—ADELAIDE, SOUTH AUSTRALIA. Lat. 34° 56′ S., Long. 138° 35′ E. Height above M.S.L. 140 Ft.

BAROMETER, WIND, EVAPORATION, LIGHTNING, CLOUDS, AND CLEAR DAYS.

	oted n. Sea Stan- ity and lings.		Wind	l.		lon		ount 9 a.m.	
Month.	Bar. corrected to 32° F. Mn. Sea Level and Stan- dard Gravity from 9 a.m. and 3 p.m. readings.	Greatest Number of Miles in one day.	Mean Hourly Pres- sure. (lbs.)	Total Miles.	Prevailing Direction.	Mean Amount of Evaporation (inches).	No. of Days Lightning.	Mean Amount of Clouds, 9 a.1 3 p.m. & 9 p.n	No. of Clear Days.
No. of yrs. over which observation extends	68 .	47	47	47	47	55	53	57	43
January February March April May June July August September October November December .	29.915 29.952 30.038 30.122 30.122 30.093 30.129 30.099 30.037 29.997 29.974 29.918	758 19/99 691 22/96 628 9/12 773 10/96 760 9/80 750 12/78 674 25/82 773 31/97 720 2/87 768 28/98 677 2/04 675 12/91	0.34 0.29 0.24 0.21 0.25 0.24 0.28 0.31 0.34 0.33 0.34	7,907 6,713 6,660 6,079 6,258 6,589 6,721 7,146 7,296 7,869 7,540 7,892	S S S S W N X E N N W W S W X W S S W S S W	8.96 7.34 5.80 3.43 2.02 1.23 1.28 1.87 2.84 4.75 6.54 8.39	2.3 2.0 2.1 1.5 1.7 2.1 1.6 2.1 2.5 3.4 3.6 2.7	3.5 3.4 3.9 5.0 5.8 6.2 5.6 5.2 4.6 3.8	8.2 7.2 7.0 4.1 1.9 1.6 1.8 2.6 3.1 3.9 5.2 7.4
	30.033	— 773 (a)	0.28	7,057	S W x S	54.45	27.6	4.8	54.0

(a) 10/4/96 and 31/8/97.

TEMPERATURE AND SUNSHINE.

		n Tem e (Fal			e Shade ire (Fah r .).	ne.	Extr Temperat	eme ure (Fahr.).	of Ine.
Month.	Mean Max.	Mean Min.	Mean.	Highest.	Lowest.	Extreme Range.	Highest in Sun.	Lowest on Grass.	Mean Hours of Sunshine.
No. of yrs. over which observation extends	68	68	68	68	68	68	47	64	43
January February March April June July August September October November December	86.3 86.2 80.7 73.3 65.6 60.3 58.9 62.0 66.2 72.4 78.6 83.3	61.6 62.1 58.8 54.6 50.3 46.7 44.6 45.9 47.9 51.4 55.3 58.9	73.9 74.1 69.8 63.9 57.9 53.5 51.7 54.0 57.1 61.9 67.0 71.1	116.3 26/58 113.6 12/99 108.0 12/61 98.0 10/66 89.5 4/21 76.0 23/65 74.0 11/06 85.0 31/11 90.7 23/82 102.9 21/22 113.5 21/65 114.2 14/76	45.1 21/84 45.5 23/18 44.8 -/57 39.6 15/59 36.9 (a) 32.5 27/76 32.0 24/08 32.3 17/59 32.7 4/58 36.0 -/57 40.8 2/09 43.0 (b)	71.2 68.1 63.2 58.4 52.6 43.5 42.0 52.7 58.0 66.9 72.7 71.2	180.0 18/82 170.5 10/00 174.0 17/83 155.0 1/83 148.2 12/79 138.8 18/79 134.5 26/90 140.0 31/92 160.5 23/82 162.0 30/21 166.9 20/78 175.7 7/99	36.5 14/79 36.7 (c) 33.8 27/80 30.2 16/17 25.9 10/91 22.9 12/13 23.3 25/11 23.5 7/88 26.2 15/08 27.8 2/18 31.5 2/09 32.5 4/84	309,5 263,8 239,7 178.0 148.1 121.0 137.6 163.7 183.1 227.2 262.4 302.4
Year {Averages Extremes	72.8	53.2	63.0	116.3 26/1/58	32.0 24/7/08	84.3	180.0 18/1/82	22.9	2536.5d

(a) 26/1895 and 24/1904. (b) 16/1861 and 4/1906.

(c) 24/78 and 23/18.

(d) Total for year.

HUMIDITY, RAINFALL, AND D

	Vapour Pressure.	Rel.	Hum.	n. (%) Rainfall (inches).									Dew.
Month.	Mean 9 a.m.	Mean 9 a.m.	Highest Mean.	Lowest Mean.	Mean Monthly.	Monthly. Mean No. of Days Rain. Greatest Monthly. Least Monthly.		Greatest	in One Day.	Mean No. Days Dew.			
No. of yrs. over which observation extends	57	57	57	57	86	86	36 86		86			86	53
January February March April May June July August September October November December	.342 .359 .348 .336 .319 .300 .276 .256 .297 .300 .312 .325	38 41 47 56 67 77 76 69 61 51 43 39	59 56 58 72 76 84 87 77 72 67 57	30 30 36 37 49 69 68 54 44 29 31	0.73 0.66 1.05 1.76 2.76 3.15 2.65 2.51 2.04 1.74 1.16	4 6 9 14 16 16 14 11 8 6	4.00 2.89 4.60 6.78 7.75 8.58 5:38 6.24 5.83 3.83 3.55 3.98	1850 1919 1878 1853 1875 1916 1865 1852 1923 1870 1851 1861	nil nil 0.03 0.20 0.42 0.37 0.35 0.45 0.17 0.04 nil	(a) (b) (c) 1923 1891 1886 1899 1914 1896 1914 1885	2.30 2.24 3.50 3.15 2.75 2.11 1.75 2.23 1.59 2.24 1.88 2.42	2/89 14/13 5/78 5/60 1/53 1/20 10/65 19/51 20/23 16/08 28/58 23/13	4.2 5.7 11.1 13.9 15.7 17.3 16.5 15.7 12.7 7.0 4.9
Totals	.311	53	=		21.21	124	-			=		_	140.4
Extremes			87	29	<u> </u>	<u> </u>	8.58	6/16	liail	(d)	3.50	5/3/78	

⁽a) 1848, &c. (b) 1848, &c.

CLIMATOLOGICAL DATA—BRISBANE, QUEENSLAND.

LAT. 27° 28' S., LONG. 153° 2' E. HEIGHT ABOVE M.S.L. 137 FT. BAROMETER, WIND, EVAPORATION, LIGHTNING, CLOUDS, AND CLEAR DAYS.

	ed tan- tan- y and ngs.		Wind			nt Ion		nnt a.m. p.m.	
Month.	Bar. corrected to 32° F. Mn. Sec Level and Stan- dard Gravity from 9 a.m. and 3 p.m. readings.	Greatest Number of Miles in one day.	Mean Hourly Pres- sure. (lbs.)	Total Miles.	Prevailing Direction.	Mean Amount of Evaporation (inches).	No. of Days Lightning.	Mean Amount of Clouds, 9 a.m. 3 p.m. & 9 p.m.	No. of Clear Days.
No. of yrs. over which observation extends.	38	14	14	14	38	16	38	33	16
January February March April May June July September October November December	29.867 29.902 29.958 30.050 30.086 30.064 30.072 30.096 30.036 30.003 29.955 29.888	361 1/22 347 5/22 305 29/16 370 27/23 307 20/22 400 12/24 339 2/23 331 6/23 322 14/23 325 25/18 274 18/23 308 24/24	0.12 0.13 0.10 0.09 0.08 0.10 0.08 0.09 0.08 0.09 0.12 0.13	4,696 4,600 4,272 3,980 3,759 3,701 3,804 4,042 3,766 4,149 4,503 4,846	E & SE SE & S	6.881 5.332 4.947 3.784 2.882 2.235 2.463 3.056 3.922 5.382 6.099 6.831	6.0 5.3 4.6 3.2 3.3 2.2 2.4 3.6 5.9 6.9 8.4 8.8	5.7 5.7 5.2 4.6 4.3 4.2 3.7 3.5 4.0 4.7 5.2	3.3 1.9 5.1 7.7 8.8 8.8 11.5 11.8 12.2 8.4 6.6 3.7
	29.998		0.10	4,176	S to E and N E	53.814	60.6	4.5	89.8

TEMPERATURE AND SUNSHINE.

		n Tem re (Fa		Extrem Temperatu		ne .	Extr Temperat	of me.	
Month.	Mean Max.	Mean Min.	Mean.	Highest.	Lowest.	Extreme Range.	Highest in Sun.	Lowest on Grass.	Mean Hours of Sunshine.
No. of yrs. over which observation extends	38	38	38	38	38	38	38	38	16
January February March April May June July August September October November December	85.5 84.5 82.4 79.0 73.7 69.4 68.4 71.1 75.8 79.8 82.9 85.2	68.9 68.3 66.3 61.6 55.4 51.0 48.6 49.8 54.9 59.9 64.2 67.5	77.2 76.4 74.4 70.3 64.6 60.2 58.5 60.4 65.4 69.8 73.6 76.4	108.9 14/02 101.9 11/04 99.4 5/19 95.2 (b) 90.3 21/23 88.9 19/18 83.4 28/98 87.5 28/07 95.2 16/12 101.4 18/93 106.1 18/13 105.9 26/93	58.8 4/93 58.7 (a) 52.4 29/13 48.6 17/00 41.3 24/99 36.3 29/08 36.1 (c) 37.4 6/87 40.7 1/96 43.3 3/99 48.5 2/05 56.4 13/12	50.1 43.2 47.0 46.6 47.5 52.6 47.3 50.1 54.5 58.1 57.6 49.5	166.4 10/17 165.2 6/10 160.0 1/87 153.8 11/16 147.0 1/10 136.0 3/18 146.1 20/15 141.9 20/17 155.5 26/03 157.4 31/18 162.3 7/89 160.4 7/14	49.9 4/93 49.3 9/89 45.4 29/13 37.0 17/00 29.8 8/97 25.4 23/88 23.9 11/90 30.4 1/89 34.9 8/89 49.1 3/94	225.4 203.8 210.1 203.7 197.5 169.9 196.1 231.2 233.5 249.9 243.2 241.1
Year { Averages Extremes	78.1	59.7	68.9	108.9 14/1/02	36.1 (c)	72.8	166.4 10/1/17	23.9	2605.4d

(a) 10 and 11/04. (b) 9/96 and 5/03. (c) 12/94 and 2/96. (d) Total for year.

HUMIDITY, RAINFALL, AND DEW.

	Vapour Pressure	Rel.	Hum.	(%)			R	ainfall (inches)) .		Dew.
Month.	Mean 9 a.m.	ean a.m.	Highest Mean.	Lowest Mean.	Mean Monthly.	Mean No. of Days Rain.	Greatest	onthly.	438	Monthly.	Greatest in One Day.	Mean No. Days Dew.
	≥ 0	₹6		HE	ZZ	2 2 H	3	Σ	1	Z	D E.C	ZΩ
No. of yrs. over which observation extends	38	38	38	38	73	65	73 27.72 1895		73		. 73	38
January February	.654 .655 .628 .533 .430 .352 .333 .348 .418 .485 .551	67 70 72 73 74 74 74 69 65 61 60 63	79 82 85 80 85 84 81 80 76 72 72 70	53 55 56 60 64 67 61 47 49 46 52	6.31 5.76 3.61 2.83 2.67 2.35 2.11 2.08 2.57 3.65 4.89	14 14 15 12 10 8 8 7 9	27.72 40.39 34.04 15.28 13.85 14.03 8.46 14.67 5.43 9.99 12.40 13.99	1895 1893 1870 1867 1876 1873 1889 1879 1886 1882 1917	0.32 0.58 nil 0.04 nil nil nil nil 0.10 0.14 nil 0.35	1919 1849 1849 1897 1846 1847 1841 (b) 1907 1900 1842 1865	18.31 21/8: 8.36 16/9: 11.18 14/06 4.47 13/16 5.62 9/7: 6.01 9/9: 3.54 (a) 4.89 12/8: 2.46 ·2/94 1.95 20/8: 4.46 16/86 6.60 28/71	6.5 9.9 12.3 13.5 11.5 12.8 10.8 10.7 8.8 5.8
Year { Totals Averages Extremes	.500	68 -	 85		45.14	128 	40.39 2/1893		nil (c)		18.31 21/1/87	114.3

(a) 15/76, 16/89. (b) 1862, 1869, 1880. (c) March, May, June, July, Aug., Nov., various years.

CLIMATOLOGICAL DATA-SYDNEY, NEW SOUTH WALES.

Lat. 33° 52′ S., Long. 151° 12′ E. Height above M.S.L., 138 Ft. Barometer, Wind, Evaporation, Lightning, Clouds, and Clear Days.

	ed tan- tan- rrly		Wind	l.		g g		ount 9 a.m., 9 p.m.	
Month.	Jar. corrected to 32° F. Mn. Se Level and Standard Gravity from 24 hourly readings.	Greatest Number of Miles in One Day.	Mean Hourly Pres- sure. (lbs.)	Total Miles.	Prevailing Direction.	Mean Amount of Evaporation (inches).	No. of Days Lightning.	Mean Amount of Clouds, 9 a.n 3 p.m. & 9 p.n	No. of Clear Days.
No. of yrs. over which observation extends	66	58	58	58	58	45	65	63	61
January February February February March April May June June June July August September Cotober November December	29.898 29.945 30.011 30.074 30.081 30.056 30.075 30.075 29.969 29.939 29.881	721 1/71 871 12/69 943 20/70 803 6/82 758 6/98 712 7/00 930 17/79 756 22/72 064 6/74 926 4/72 720 13/68 938 3/84	0.33 0.31 0.24 0.21 0.22 0.27 0.27 0.25 0.29 0.31 0.32	8,066 6,921 6,714 6,065 6,281 6,834 7,033 6,790 7,040 7,688 7,523 7,937	NENENEWWWWWWWWNENE	5.292 4.145 3.576 2.556 1.774 1.426 1.512 1.877 2.654 3.836 4.565 5.332	4.8 4.2 4.1 3.8 3.3 2.2 2.4 3.2 4.1 4.9 5.4	5.8 6.0 5.5 5.1 4.8 4.4 4.4 4.9 5.5 5.6	2.4 1.7 2.3 3.2 3.8 4.2 5.0 5.4 4.5 3.0 2.2 2.4
Year { Totals Averages Extremes	30.000	964 6/9/74	0.28	7,074	N E	38.545	48.1	5.1	40.1

TEMPERATURE AND SUNSHINE.

XI43-		Temp e (Fah			e Shade ire (Fahr.).	eg .		Extreme Temperature (Fahr.).		
Month.	Mean Max.	Mean Min.	Mean.	Highest.	Lowest.	Extreme Range.	Highest in Sun.	Lowest on Grass.	Mean Hours of Sunshine.	
No. of yrs. over which observation extends	66	66	66	66	66	66	63	66	14	
January February February March April May June July August September October November December	78.5 77.6 75.7 71.2 65.4 60.9 59.4 62.6 66.9 71.2 74.5 77.3	64.9 64.9 62.9 58.1 52.2 48.2 45.9 47.5 51.5 55.8 59.6 62.9	71.7 71.2 69.3 64.6 58.8 54.6 52.6 55.0 59.2 63.5 67.1 70.1	108.5 13/96 101.9 18/23 102.6 3/69 91.0 20/22 86.0 1/19 79.8 2/23 74.9 17/71 82.0 31/84 92.3 27/19 99.7 19/98 102.7 21/78 107.5 31/04	51.2 14/65 49.3 28/63 48.8 14/86 44.6 27/64 40.2 22/59 38.0 5/20 35.9 12/90 36.8 3/72 40.8 18/64 42.3 3/18 45.8 1/05 48.4 3/24	57.3 52.6 53.8 46.4 45.8 41.8 39.0 45.2 51.5 57.4 56.9 59.1	164.3 26/15 156.5 7/64 158.0 19/11 144.1 10/77 129.7 1/96 125.5 2/23 124.7 19/77 149.0 30/78 149.0 30/78 142.2 12/78 151.9 (a) 158.5 28/99 164.5 27/89	44.2 18/97 43.4 25/91 39.9 17/13 33.3 24/09 29.3 25/17 28.1 24/11 24.0 4/93 30.1 17/05 32.7 9/05 36.0 6/08 41.4 3/24	210.0 182.2 193.1 151.4 138.1 125.8 135.8 185.8 187.6 208.0 200.4 203.8	
Year { Averages Extremes	70.1	56.2	d 3.2	108.5 35.9 72.6 164.5 24.0		24.0 4/7/93	2125.0 b			

					, -,	,		· 		
	Vapour Pressure (inches.)	Rel.	Hum.	(%)			Rainfal	l (inches).		Dew.
Month.	Mean 9 a.m.	Mean 9 a.m.	Highest Mean.	Lowest Mean.	Mean Monthly.	Mean No. of Days Rain.	Greatest Monthly.	Least Monthly.	Greatest in One Day.	Mean No. Days Dew.
No. of yrs. over which observation extends		66	66	66	66	66	66	66	66	66
January February March April May June July August September October November December	.556 .567 .534 .459 .335 .292 .269 .352 .402 .466 .530	67 70 72 76 74 76 75 71 67 63 64 66	78 81 85 87 90 89 88 84 79 77	58 59 62 63 63 68 65 56 49 46 42 52	3.65 4.33 4.92 5.04 4.78 4.97 3.02 2.90 2.91 2.82 2.82	14 14 15 14 15 13 12 11 12 12 12 12	15.26 191 18.56 187 18.70 187 24.49 186 23.03 191 16.30 188 13.21 190 14.89 189 14.05 187 11.14 191 9.89 186 15.82 192	3	7.08 13/11 8.90 25/73 6.52 9/13 7.52 29/60 8.36 28/89 5.17 16/84 5.72 28/08 5.33 2/60 5.69 10/79 6.37 13/02 4.23 19/00 4.75 13/10	1.2 2.0 3.3 5.5 6.2 5.3 5.3 4.9 3.4 3.0 2.1
Year { Totals	.421	70	90	_ 	47.58 —	157	24.49 April/6	0.04 Aug./85	8.90 25/2/73	43.6

CLIMATOLOGICAL DATA-MELBOURNE, VICTORIA.

Lat. 37° 49′ S., Long. 144° 58′ E. Height above M.S.L., 115 Ft. BAROMETER, WIND, EVAPORATION, LIGHTNING, CLOUDS, AND CLEAR DAYS.

	rected Mn. Sea d Stan- avity m., 3 & adings.		Wind	1.		B st		ount 9a.m., 9 p.m.	
Month.	Bar. corrected to 32° F. Mn. Sca Level and Standard Gravity from 9 a.m., 3 & 9 p.m. readings.	Greatest Number of Miles in One Day.	Mean Hourly Pres- sure. (lbs.)	Total Miles.	Prevailing Direction.	Mean Amount of Evaporation (inches).	No. of Days Lightning.	Mean Amount of Clouds, 9 a.n 3 p.m. & 9 p.n	No. of Clear Days.
No. of yrs. over which observation extends	67	51	51	51	51	52	17	67	17
January February March April May June July August September October November December	29.910 29.960 30.033 30.105 30.107 30.094 30.065 29.996 29.965 29.948 29.897	583 10/97 566 8/88 677 9/81 597 7/88 693 12/65 761 13/76 637 14/75 617 11/72 899 5/66 655 1/75	0.27 0.25 0.20 0.19 0.19 0.22 0.21 0.24 0.26 0.27 0.27	7,269 6,308 6,275 5,666 5,849 6,345 6,762 6,762 6,743 7,231 6,953 7,393	S W, S E S W, S E S W, N W N W, N E N W, N E N W, N E N W, S W S W, S E S W, S E	6.426 5.056 3.976 2.375 1.469 1.094 1.050 1.468 2.299 3.341 4.538 5.761	1.8 2.7 1.6 0.8 0.6 0.8 0.5 1.1 1.6 2.2 2.5 1.9	5.1 5.0 5.5 5.8 6.5 6.7 6.3 6.3 6.1 6.0 5.9	7.4 7.1 5.1 4.6 3.2 2.2 3.2 2.9 3.6 4.1 3.5 4.2
\mathbf{Y} ear $\left\{egin{array}{ll} \mathbf{Totals} & \dots \\ \mathbf{A}\mathbf{verages} & \dots \\ \mathbf{E}\mathbf{xtremes} & \dots \end{array}\right.$	30.012	— 899 5/10/66	0.24	6,608	s w, n w	38.853	18.1	5.9	51.1 —

TEMPERATURE AND SUNSHINE.

,		Tempera- (Fahr.).		e Shade ire (Fahr.).	ne.	Extr Temperati	eme ure (Fahr.).	of Ine.
Month.	Mean M Max. M	Iean Iin. Mear	Highest.	Lowest.	Extreme Range.	Highest in Sun.	Lowest on Grass.	Mean Hours of Sunshine.
No. of yrs. over which observation extends		69 69	69	69	69	65	65	43
January February February March April May June July August September October November December	78.0 5 74.3 5 68.4 5 61.5 4 56.8 4 55.6 4 58.7 4 62.6 4 67.0 4 71.3 5	56.8 67.4 67.5 54.6 64.5 54.6 64.5 50.7 59.5 46.7 54.1 44.1 50.4 411.7 48.6 43.4 51.0 48.2 57.6 51.2 61.3 64.8	109.5 7/01 105.5 2/93 94.0 (a) 83.7 7/05 72.2 1/07 68.4 24/78 77.0 20/85 85.0 19/19 98.4 24/14 105.7 27/94	42.0 28/85 40.2 24/24 37.1 17/84 34.8 24/88 29.9 29/16 28.0 11/60 27.0 21/69 28.3 11/63 31.1 16/08 32.1 3/71 36.5 2/96 40.0 4/70	69.2 69.3 68.4 59.2 53.8 44.2 41.4 48.7 53.9 66.3 69.2 70.7	178.5 14/62 167.5 15/70 164.5 1/68 152.0 8/61 142.6 2/59 129.0 11/61 125.8 27/80 137.4 29/69 142.1 20/67 154.3 28/68 159.6 29/65 170.3 20/69	30.2 28/85 30.9 6/91 28.9 (h) 25.0 23/97 21.1 26/16 20.4 17/95 20.5 12/03 21.3 14/02 22.8 8/18 24.8 22/18 24.6 2/96 33.2 1/04	266.8 244.4 207.3 163.1 141.2 111.5 106.8 155.5 172.1 206.7 244.2 256.9
	67.3 4	19.5 58.4	111.2 14/1/62	27.0 21/7/69	84.2	178.5 14/1/62	20.4	2277.0c
Extremes	865 and	- l 17/1922	14/1/62		<u> </u>		17/6/95	_

	Vapour Pressure	Rel.	Hum.	(%)]	Rainfall	(inches).			Dew.
Month.	Mean 9 a m.	Mean 9 a.m.	Highest Mean.	Lowest Mean.	Mean Monthly.	Mean No. of Days Rain.	4 % 8 		Greatest Monthly. Least Monthly.		Greatest	in One Day.	Mean No. Days Dew
No. of yrs. over which observation extends	17	17	17	17	69	69		39		39		66	17
January February March April May June July August September October November December	.383 .417 .376 .338 .318 .283 .260 .272 .292 .304 .330 .358	58 62 64 71 78 83 82 76 69 62 59	65 69 71 78 84 88 86 82 76 67 69	50 53 57 66 71 77 76 70 60 56 52 51	1.87 1.74 2.23 2.21 2.18 2.08 1.85 1.89 2.47 2.67 2.26 2.33	8 7 9 11 13 14 14 14 14 13 11	5.68 6.24 7.50 6.71 4.31 4.51 7.02 4.04 7.93 7.61 6.71 7.18	1904 1904 1911 1901 1862 1859 1891 1924 1916 1869 1916	0.04 0.03 0.18 Nil 0.45 0.73 0.57 0.48 0.52 0.29 0.25 0.11	1878 1870 1859 1923 1901 1877 1902 1903 1907 1914 1895 1904	2.97 3.37 3.55 2.28 1.85 1.74 2.71 1.94 2.62 3.00 2.57 2.62	9/97 18/19 5/19 22/01 7/91 21/04 12/91 26/24 12/80 17/69 16/76 28/07	2.1 2.7 7.2 8.4 8.6 8.3 10.0 8.3 6.4 6.2 1.9 2.0
Year { Totals Averages Extremes	.319	68	 88		25.78	137	7.93	9/16	Nil	4/23	3.55	5/3/19	72.1

CLIMATOLOGICAL DATA-HOBART, TASMANIA.

Lat. 42° 53′ S., Long. 147° 20′ E. Height above M.S.L., 177 Ft. BAROMETER, WIND, EVAPORATION, LIGHTNING, CLOUDS, AND CLEAR DAYS.

	oted n. Sea Stan- lty i. and lings.		W	# B		a.m.			
Month.	Bar. corrected to 32° F. Mu. Sea Level and Stan- dard Gravity from 9 a.m. and 3 p.m. readings.	Greatest Number of Miles in One Day.	Mean Hourly Pres- sure. (lbs.)	Total Miles.	Prevailing Direction. 9 a.m. 3 p.m	Mean Amount of Evaporation (inches).	No. of Days Lightning.	Mean Amount of Cloud, 9 a.m 3 p.m. & 9 p.n	No. of Clear Days.
No. of yrs. over which observation extends	40	14	14	14	19	14	17	62	18
January February March April May June July August September October November December	29.826 - 29.925 29.937 29.965 29.982 29.938 29.934 29.927 29.814 29.748 29.748	500 30/16 393 19/13 407 16/21 475 12/22 411 3/16 569 27/20 425 16/21 459 30/11 516 26/15 461 8/12 508 18/15	0.19 0.13 0.13 0.14 0.12 0.13 0.12 0.13 0.18 0.18 0.20 0.18	5,995 4,567 4,962 4,877 4,844 4,757 4,767 4,767 5,514 5,757 5,878 5,725	NNW SE	5.045 3.714 2.967 1.990 0.891 0.899 1.247 1.985 3.116 4.030 4.562	0.9 1.4 1.2 0.8 0.6 0.7 0.6 0.9 0.9 0.9	5.9 6.0 5.9 6.0 6.1 5.8 5.9 6.1 6.3 6.3	2.5 2.5 2.0 1 8 2.1 1.7 2.6 2.1 1.7 1.6 1.6
	29.893		0.15	5,210	NNW SE&	31.825	10.6	6.0	23.3

TEMPERATURE AND SUNSHINE.

	Mean Tempera- ture (Fahr.).				e Shade ire (Fahr.).	e e	Extr Temperat	of ne.	
Month.	Mean Max.	Mean Min.	Mean.	Highest.	Lowest.	Extreme Range.	Highest in Sun.	Lowest on Grass.	Mean Hours of Sunshine.
No. of yrs. over which observation extends January February March April May June July August September October November December Vear	71.3 71.3 68.0 62.7 57.4 52.8 52.0 55.0 58.8 62.7 66.1 69.4	54 52.9 53.3 50.8 47.6 43.6 41.0 39.3 41.1 43.2 45.4 48.3 51.2	54 62.1 62.3 59.4 55.2 50.5 46.9 45.6 48.0 51.0 54.0 57.2 60.3	78 105.0 1/00 104.4 12/99 98.8 5/46 90.0 2/56 77.8 5/21 75.0 7/74 72.0 22/77 77.0 3/76 80.0 9/72 92.0 24/14 98.0 20/88 105.2 30/97	78 40.3 (a) 39.0 20/87 36.0 31/05 39.0 25/56 39.2 20/02 28.0 22/79 27.0 18/66 30.0 12/41 32.0 12/89 35.2 5/13 38.0 13/06	78 64.7 65.4 62.8 60.0 48.6 47.0 45.0 60.0 62.8 67.2	37 160.0 (b) 165.0 24/98 150.0 3/05 142.0 18/93 128.0 (c) 122.0 12/94 121.0 12/93 129.0 -/87 138.0 23/93 156.0 9/93 154.0 19/92 157.0 30/18	57 30.6 19/97 28.3 -/87 27.5 30/02 25.0 -/86 20.0 19/02 21.0 6/87 18.7 16/86 20.1 7/09 22.3 20/14 23.8 (d) 26.0 1/08 27.2 -/86	30 210.2 177.2 173.0 140.5 132.3 102.9 124.1 144.2 144.5 168.9 198.1 194.1
(a) 3/72 and 2/06.	(b) 5	/86 an	d 13/0	30/12/97	18/7/66 and -/92.		24/2/98 /86 and -/99.	16/7/86 (e) Total for	Venr

	Vapour Prossure (inches)	Rel. Hum. (%)		Rainfall (inches).									
Month.	Mean 9 a.m.	Mean 9 a.m.	Highest Mean.	Lowest Mean.	Mean Monthly.	Mean No. of Days Rain.	Greatest Monthly.		Least Monthly.		Greatest in One Day.		Mean No. Days Dew.
No. of yrs. over which observation extends	41	41	41	41	82	81	82		82		58		15
January February March April May June July August September October November December	.348 .356 .323 .307 .267 .245 .234 .249 .266 .273 .295 .312	63 65 68 73 78 82 80 77 71 66 63 61	77 80 78 84 88 92 88 85 82 80 78	51 51 58 61 68 68 72 64 60 51 50	1.85 1.46 1.71 1.86 1.87 2.21 2.17 1.84 2.11 2.24 2.52 1.99	10 8 10 12 13 14 14 14 14 15 14	5.91 9.15 7.60 6.50 6.37 8.15 6.02 10.16 7.14 6.67 8.92 9.00	1893 1854 1854 1909 1905 1889 1922 1858 1844 1906 1849 1875	0.03 0.07 0.02 0.07 0.10 0.22 0.30 0.23 0.39 0.26 0.16	1841 1847 1843 1904 1843 1852 1850 1854 1847 1850 1868 1842	2.96 4.50 2.79 5.02 3.22 4.11 2.51 4.35 3.50 2.58 3.97 2.48	30/16 25/54a 5/19 20/09 14/58 14/89 18/22 12/58 29/44 4/06 6/49 13/16	0.7 1.9 4.4 10.6 13.3 8.4 8.2 9.0 4.7 3.1 1.3 0.9
	.282	70	92	49	23.83	149	— 10.16 8/1858		0.02		5.02 20/4/09		66.5