

CHAPTER II.

PHYSIOGRAPHY.

§ 1. General Description of Australia.

1. *Geographical Position.*—(i) *General.* The Australian Commonwealth, which includes the island continent of Australia proper and the island of Tasmania, is situated in the Southern Hemisphere, and comprises in all an area of about 2,974,581 square miles, the mainland alone containing about 2,948,366 square miles. Bounded on the west and east by the Indian and Pacific Oceans respectively, it lies between longitudes 113° 9' E. and 153° 39' E., while its northern and southern limits are the parallels of latitude 10° 41' S. and 43° 39' S., or, excluding Tasmania, 39° 8' S. On its north are the Timor and Arafura Seas and Torres Strait, on its south the Southern Ocean. The extreme points are Steep Point on the west, Cape Byron on the east, Cape York on the north, and South-East Cape or, if Tasmania be excluded, Wilson's Promontory, on the south.

(ii) *Tropical and Temperate Regions.* Of the total area of Australia, nearly 39 per cent. lies within the tropics. Assuming, as is usual, that the latitude of the Tropic of Capricorn is 23° 30' S., the areas within the tropical and temperate zones are approximately as follows:—

AUSTRALIA: AREAS OF TROPICAL AND TEMPERATE REGIONS.

(Square miles.)

Area.	N.S.W. (a)	Vic.	Qld.	S. Aust.	W. Aust.	Tas.	N. Terr.	Total.
Within Tropical Zone ..			359,000		364,000		426,320	1,149,320
„ Temperate Zone	310,372	87,884	311,500	380,070	611,920	26,215	97,300	1,825,261
Total Area ..	310,372	87,884	670,500	380,070	975,920	26,215	523,620	2,974,581

(a) Includes Australian Capital Territory (939 square miles).

Fifty-four per cent. of Queensland lies within the tropical zone and 46 per cent. in the temperate zone; 37 per cent. of Western Australia is tropical and 63 per cent. temperate; while 81 per cent. of the Northern Territory is tropical and 19 per cent. temperate. All of the remaining States lie within the temperate zone. The tropical part of Australia thus comprises about 39 per cent. of the whole of the continent, and about 53 per cent. of the three territories which have areas within the tropical zone.

2. *Area of Australia compared with Areas of other Countries.*—The area of Australia is almost as great as that of the United States of America, four-fifths of that of Canada, nearly three-quarters of the whole area of Europe, and about 25 times that of Great Britain and Ireland. The areas of Australia and of certain other countries are shown in the table on the following page.

AREA OF AUSTRALIA AND OF OTHER COUNTRIES, *circa* 1956.

('000 square miles.)

Country.	Area.	Country.	Area.
Continental Divisions—		Africa—continued.	
Europe (a)	1,903	Rhodesia and Nyasaland Federation	488
Asia (a)	10,494	Angola	481
U.S.S.R. (Europe and Asia)	8,650	Union of South Africa	472
Africa	11,695	Ethiopia and Eritrea	457
North and Central America and West Indies	9,354	Egypt	386
South America	6,856	Nigeria and Protectorate (b)	373
Oceania	3,304	Tanganyika Territory	363
<i>Total, excluding Arctic and Antarctic Conts. ..</i>	<i>52,256</i>	South-West Africa	318
Europe (a)—		Mozambique	298
France	213	Bechuanaland Protectorate	275
Spain (incl. possessions) ..	194	Madagascar	228
Sweden	174	Kenya Colony and Protectorate	225
Germany	137	Other	1,100
Finland	130	<i>Total</i>	<i>11,695</i>
Norway	125	North and Central America—	
Poland	120	Canada	3,851
Italy	116	United States of America ..	3,022
Yugoslavia	99	Alaska	586
United Kingdom	94	Greenland	840
Romania	92	Mexico	760
Other	409	Nicaragua	57
<i>Total (a)</i>	<i>1,903</i>	Cuba	44
Asia (a)—		Honduras	43
China, Mainland	3,769	Other	151
India and Nepal	1,270	<i>Total</i>	<i>9,354</i>
Iran	629	South America—	
Saudi Arabia	618	Brazil	3,287
Mongolian People's Republic	591	Argentina	1,073
Indonesia	576	Peru	482
Pakistan	365	Colombia (excl. of Panama)	440
Turkey	300	Bolivia	424
Burma	262	Venezuela	352
Afghanistan	251	Chile	286
Thailand	198	Paraguay	157
Iraq	172	Ecuador	105
Other	1,493	Other	250
<i>Total (a)</i>	<i>10,494</i>	<i>Total</i>	<i>6,856</i>
U.S.S.R.	8,650	Oceania—	
Africa—		Commonwealth of Australia	2,975
French West Africa	1,789	New Zealand	104
French Equatorial Africa ..	969	New Guinea (c)	93
Sudan	968	Papua	91
Algeria	920	Other	41
Belgian Congo	906	<i>Total</i>	<i>3,304</i>
Libya	679		

(a) Excludes U.S.S.R., shown below. (b) Includes British Cameroons. (c) Australian Trust Territory.

The areas shown in the table are obtained from the *Demographic Yearbook, 1957*, published by the Statistical Office of the United Nations and the countries have been arranged in accordance with the continental groups used therein.

3. Areas of States and Territories, Coastal Configuration and Standard Times.—As already stated, Australia consists of six States and the Northern and Australian Capital Territories. Particulars of areas, coastline and standard times are shown in the following table:—

AUSTRALIA: AREAS OF STATES AND TERRITORIES, COASTLINE AND STANDARD TIMES.

State or Territory.	Area.	Proportion of Total Area.	Coastline.	Area per Mile of Coastline.	Standard Times.	
					Meridian Selected.	Ahead of G.M.T.
	Sq. miles.	%	Miles.	Sq. miles.		Hours.
New South Wales ..	309,433	10.40	(a) 700	(a) 443	150° E.	10
Victoria ..	87,884	2.96	680	129	150° E.	10
Queensland ..	670,500	22.54	3,000	223	150° E.	10
South Australia ..	380,070	12.78	1,540	247	142°30'E.	9½
Western Australia ..	975,920	32.81	4,350	224	120° E.	8
Northern Territory ..	523,620	17.60	1,040	503	142°30'E.	9½
Australian Capital Territory ..	939	0.03	150° E.	10
<i>Mainland</i>	<i>2,948,366</i>	<i>99.12</i>	<i>11,310</i>	<i>261</i>
Tasmania	26,215	0.88	900	29	150° E.	10
Australia ..	2,974,581	100.00	12,210	244

(a) Includes Australian Capital Territory.

There are few striking features in the configuration of the coast; the most remarkable indentations are the Gulf of Carpentaria on the north and the Great Australian Bight on the south. The Cape York Peninsula on the extreme north is the only other remarkable feature in the outline. In Official Year Book No. 1 an enumeration was given of the features of the coastline of Australia (*see* pages 60–68).

Prior to 1895, the official time adopted in the several colonies was for most purposes the mean solar time of the capital city of each. In 1894 and 1895, after several conferences had been held, legislation was enacted by each of the colonies whereby the mean solar times of the meridians of east longitude 120° (Western Australia), 135° (South Australia and Northern Territory) and 150° (Queensland, New South Wales, Victoria and Tasmania) were adopted. In 1898, however, the South Australian legislature amended its earlier provision and adopted the mean solar time of the meridian 142° 30' E. longitude as the standard time for that colony. For further information on this subject, *see* Official Year Book No. 39, page 65.

4. Geographical Features of Australia.—(i) *General*. The following description is only a broad summarization of the main physical characteristics of the Australian continent. For greater detail of particular geographical elements, earlier issues of the Official Year Book should be consulted. The list of special articles, etc., at the end of this volume indicates the nature of the information available and its position in the various issues.

(ii) *Orography of Australia*. (a) *General Description of the Surface*. Owing to the absence of any very high mountain chains, and to the great depression in the centre of Australia, the average elevation of the Australian continent above the level of the surrounding oceans is less than that of any of the other continents. Three-quarters of the land-mass lies between the 600 and 1,500 feet contours in the form of a huge plateau.

A section through the continent from east to west, at the point of its greatest breadth, shows first a narrow belt of coastal plain. This plain, extending north and south along the whole eastern coast, is well watered by rivers. It is of variable width, seldom more than sixty or seventy miles, and occasionally only a few miles, the average being roughly about forty to fifty miles. From this plain, the Great Dividing Range, extending from the north of Queensland to the south of New South Wales, and thence sweeping westward through Victoria, rises, often abruptly, and frequently presents bold escarpments on its eastern face. The descent on its western slopes is gradual, until, in the country to the north of Spencer's Gulf, the plain is not above sea-level and occasionally even below it. Thence there is another almost imperceptible rise until the mountain ranges of Western Australia are reached, and beyond these lies another coastal plain.

The great central plain or plateau is the most distinctive feature of the Australian continent and its climatic peculiarities can probably be largely ascribed thereto.

(b) *Mountain Systems.* The main mountain feature of Australia is the Great Dividing Range, which runs along the whole eastern coast of the continent and can be traced over the islands of Torres Strait to New Guinea, while, in the south, one branch sweeps westwards towards the boundary of Victoria and South Australia, and the other, the main branch, terminates in Tasmania.

This mountain system is at no place more than 250 miles from the eastern coastline and it approaches to less than 30 miles. On the whole, it is much closer to the coast in both New South Wales and Victoria than it is in Queensland, the corresponding average distances being about 70, 65 and 130 miles respectively. There is no connexion between the mountains of the eastern and other States of Australia.

The mountains of Australia are relatively low. Thus, in Queensland, the Great Dividing Range reaches a height above sea-level of less than 5,500 feet, the highest peak being Mount Bartle Frere. Mount Kosciusko, in New South Wales, is only about 7,300 feet, and Mount Bogong, in Victoria, about 6,500 feet high. In South Australia and Western Australia, heights of three and four thousand feet are attained. In Tasmania the greatest height is only a little more than 5,000 feet. The fact that there are no high mountains in Australia is also of considerable importance in considering the climate of Australia.

It may be of interest to observe that at one time Tasmania was probably connected with the mainland. As the Great Dividing Range can, in the north, be traced from Cape York across Torres Strait to New Guinea, so its main axis can be similarly followed across the shallow waters of Bass Strait and its islands from Wilson's Promontory to Tasmania, which may be said to be completely occupied by ramifications of the chain. The central part of the island is occupied by an elevated plateau, roughly triangular in shape, and presenting bold fronts to the east, west and north. This does not extend in any direction more than about 60 miles. The plateau rests upon a more extensive tableland, the contour of which closely follows the coastline, and occasionally broadens out into low-lying tracts not much above sea-level. The extreme south of the island is rugged in character.

(iii) *Hydrology of Australia.* (a) *Rainfall.* On the whole, Australia is a country with a limited rainfall. This is immediately evident on studying its river systems, its lakes, and its artesian areas. Its one large river system is that of the Murray and Darling Rivers, of which the former stream is the larger and more important. Many of the rivers of the interior run only after heavy rains. Depending almost entirely on rainfall, a consequence of the absence of high mountains, they drain large areas with widely varying relation as between rainfall and flow. Thus it has been estimated that not more than 10 per cent. of the rainfall on the catchment area of the Darling River above Bourke (New South Wales) discharges itself past that town. The rate of fall is often very slight.

(b) *Rivers.* The rivers of Australia may be divided into two major classes, those of the coastal plains with moderate rates of fall; and those of the central plains with very slight fall. Of the former not many are navigable for any distance from their mouths, and bars make many of them difficult of access or inaccessible from the sea.

The two longest rivers of the northern part of the eastern coast are the Burdekin, discharging into Upstart Bay, with a catchment area of 53,500 square miles, and the Fitzroy, which reaches the sea at Keppel Bay and drains about 55,600 square miles.

The Hunter is the largest coastal river of New South Wales, draining about 11,000 square miles before it reaches the sea at Newcastle. The Murray River, with its great tributary the Darling, drains a considerable part of Queensland, the major part of New South Wales and a large part of Victoria. It debouches into the arm of the sea known as Lake Alexandrina, on the eastern side of the South Australian coast. The total length of the Murray is about 1,600 miles, 400 being in South Australia and 1,200 constituting the boundary between New South Wales and Victoria. The total length of the Darling-Murray from the source of the Darling to the mouth of the Murray is about 2,300 miles. In good seasons the river is navigable for a considerable proportion of its length.

The rivers of the north-west coast of Australia (Western Australia) are of considerable size, e.g., the Murchison, Gascoyne, Ashburton, Fortesque, De Grey, Fitzroy, Drysdale and Ord. So also are those in the Northern Territory, e.g., the Victoria and Daly. The former of these, estimated to drain 90,000 square miles, is said to be navigable for 50 miles.

The rivers on the Queensland side of the Gulf of Carpentaria, such as the Gregory, Leichhardt, Cloncurry, Gilbert and Mitchell, are also of considerable size.

Owing to the small amount of fall of many of the interior rivers, they may flood hundreds of miles of country in wet seasons, while in dry seasons they form a mere succession of waterholes or are entirely dry. It is this fact that explains the apparently conflicting reports of the early explorers, one regarding the interior as an inland sea, and another as a desert.

The rivers of Tasmania have short and rapid courses, as might be expected from the configuration of the territory.

(c) *Lakes.* The "lakes" of Australia may be divided into three classes, true permanent lakes; lakes which, being very shallow, become mere morasses in dry seasons or even dry up and finally present a cracked surface of salt and dry mud; and lakes which are really inlets of the ocean, opening out into a lake-like expanse.

The second class is the only one which seems to demand special mention. These are a characteristic of the great central plain of Australia. Some of them, such as Lakes Torrens, Gairdner, Eyre and Frome, are of considerable extent.

(d) *Artesian Areas.* A considerable tract of the plain country of New South Wales and Queensland carries a water-bearing stratum, usually at a great depth. A large number of artesian bores have been put down, from which there is a considerable flow. These are of great value and render usable large areas which otherwise would be difficult to occupy even for pastoral purposes.

For further information on this subject, see Chapter VIII.—Water Conservation and Irrigation.

5. *Fauna, Flora, Geology and Seismology of Australia.*—Special articles dealing with these features have appeared in previous issues of the Official Year Book, but limits of space preclude their repetition in each volume. The nature and location of these articles can be readily ascertained from the special index preceding the general index at the end of this volume.

§ 2. Climate and Meteorology of Australia.

NOTE.—This Section has been prepared by the Director, Commonwealth Meteorological Bureau, and the various States and Territories have been arranged in the standard order adopted by that Bureau. A special article dealing with droughts in Australia has been included in this issue (see para. 9, page 51).

1. *Introductory.*—Previous issues of the Official Year Book, notably No. 3, pages 79–83, and No. 4, pages 84 and 87, contained outlines of the history of Australian meteorology and the creation and organization of the Commonwealth Bureau of Meteorology. Official Year Book No. 38, pages 30–32, contained paragraphs devoted to (i) Organization of the Meteorological Service; (ii) Meteorological Publications; (iii) Equipment; and (iv) Meteorological Divisions.

By reason of its insular geographical position and the absence of striking physical features, whether in marine gulfs or in important mountains, Australia is, on the whole, less subject to extremes of weather than are regions of similar area in other parts of the globe, and latitude for latitude Australia is generally more temperate.

The average elevation of the surface of the land is low, probably close to 900 feet above the sea. The altitudes range up to a little more than 7,300 feet, hence the Australian climate displays a great many features, from the characteristically tropical to what is essentially alpine, a fact indicated in some measure by the name Australian Alps given to the southern portion of the Great Dividing Range.

On the coast, the rainfall is often abundant and the atmosphere moist, but in some portions of the interior it is very limited, and the atmosphere dry. The distribution of forest, therefore, with its climatic influence, is very uneven. In the interior, in places, there are fine belts of trees, but there are also large areas which are treeless, and here the air is hot and parching in summer. Again, on the coast, even so far south as latitude 35°, the vegetation is tropical in its luxuriance, and to some extent also in character.

2. *Temperature.*—(i) *Effective Temperature.* When a meteorologist speaks of temperature, he means the temperature of the air indicated by a thermometer sheltered from precipitation, from direct rays of the sun and from radiation of heat from the ground

and neighbouring objects, yet freely exposed to the circulation of the air. In other words, he means temperature measured under conditions standardized as nearly as possible in a Stevenson Screen, which is the standard housing for meteorological thermometers.

This shade temperature as measured by a "dry bulb" thermometer shows only the actual temperature experienced by dry inorganic substances, not the *sensible* temperatures felt by organic bodies. In the case of human beings, *sensible* temperature is affected by the rate of conduction of heat to or from the body by moving air and also by the rate of cooling due to evaporation from the skin and respiratory passages. The wind and humidity therefore determine the *sensible* temperature.

The humidity (relative humidity) is determined from the readings of the dry and wet bulb thermometers. Of late years, however, with increasing interest in human comfort in tropical climates, another term, *effective* temperature, has come into use. It may be defined as "the temperature of a still, saturated atmosphere which would on the average produce the same feeling of warmth or cold as the atmosphere in question".*

The 80° F. isotherm is confined to a very narrow tract of country on the north-west coast of Western Australia. The 75° F. isotherm extends, broadly, from Onslow on the north-west coast of Western Australia to Daly Waters to Camooweal to Moreton in Cape York Peninsula following in a general way the coastline of Northern Australia but from 100 to 300 miles inland.

Later investigations have established "comfort zones"† bounded by limits of effective temperature within which people will feel comfortable. American research workers have determined the following figures ‡:—

COMFORT ZONES: EFFECTIVE TEMPERATURES.

Season.	No subjects feel comfortable below—	Fifty per cent. of subjects feel comfortable between—	No subjects feel comfortable above—
Winter.. ..	60° F.	63° and 71° F.	74° F.
Summer	64° F.	66° and 75° F.	79° F.

Queensland investigators§ in recent years have divided some towns of Queensland into three classes on the basis of deviation from comfort:—

Class 1 (Sub-tropics).—Quite suitable for Caucasian habitation—Rockhampton, Bundaberg, Brisbane, Longreach, Charleville.

Class 2 (Marginal tropics).—Suitable for Caucasian habitation, but requires adaptation in summer—Mackay, Townsville.

Class 3 (Tropics).—(a) Permissible for Caucasian habitation but requires selection and marked adaptation—Cardwell, Cairns, Cloncurry. (b) Not suitable for continuous Caucasian habitation—Cape York, Burketown.

These results of recent years bear out investigations made previously in Australia|| in which the atmospheric vapour pressure was used as a measure of comfort, its value for this purpose being that it has equal effect in both indoor and outdoor climates. The limits of comfort range from 0.2 to 0.5 inch of vapour pressure. After drawing isopleths for effective temperature (not corrected for altitude), mean vapour pressure reduced to a logarithmic scale, and mean wet bulb temperature, it is found that there is close agreement in defining zones of relative discomfort.

(ii) *Seasons.* The Australian seasons are:—Summer, December to February; autumn, March to May; winter, June to August; spring, September to November. In most parts of Australia, January is the hottest month, but in Tasmania and southern Victoria, February is the hottest; in the tropical north, probably because the cooling "monsoon" rains occur in late summer, December is the hottest month, and at Darwin, November.

On a rainfall basis, in the tropical north the year is divisible into "wet" and "dry" seasons, but on the basis of temperatures and physical comfort the "dry" season can be further sub-divided into two parts—"cool dry" and "warm dusty".¶

(a) "*Cool dry*" Season. From May to August. The average maximum temperature ranges from 80° to 85° F., the relative humidity is low and in inland areas cold nights are experienced when the temperature drops to 40° F. The skies generally are cloudless, but in about one year in three during June or July one to two inches of rain fall.

* Houghton, F. C., Teague, W. W. and Miller, W. E. (1926) Amer. Soc. Heat. Vent. Engrs. † Yaglou, C. P. (1926) J. Industr. Hyg. ‡ Yaglou, C. P. (1927) Ibid. § Lee, D. H. K. Trans. Roy. Soc. Trop. Med. and Hyg. (1940) Vol. XXXII. ¶ Barkley, H. Zones of Relative Physical Comfort in Australia, Met. Bull. 20, 1934. ¶ Maze, W. H. Austr. Geog. June, 1945. Settlement in E. Kimberleys.

(b) "*Warm dusty*" Season. From the end of August, temperatures rise and reach a maximum in October or the beginning of November. Temperatures of over 120° F. have been recorded.

(c) "*Wet*" Season. After the first of the heavy storms, the maximum temperatures fall but still remain high with high relative humidity. At Wyndham during January, 1944, the minimum temperature did not drop below 75° F. for fourteen consecutive days. A maximum of over 100° F. was recorded on each rainless day.

In Central and Northern Australia, during the hottest months, the average temperatures range from 80° to 85° F., whereas in Southern Australia they vary from 65° to 70° F. (*see* maps pages 33, 34).

Throughout Australia, the coldest month is July, when only a very narrow strip of the northern sea-board has an average temperature as high as 75° F. Over the southern half of the continent, July temperatures range from 55° to 45° F. at elevations below 1,500 feet and fall as low as 35° on the Australian Alps (*see* maps pages 35, 36). Here the temperature seldom, if ever, reaches 100° F. even in the hottest of seasons. Hotham Heights (6,100 feet above Mean Sea Level) recorded the highest maximum of 82.0° F. on 20th January, 1935. In winter, readings slightly below zero are occasionally recorded on the extreme heights.

Tasmania, as a whole, enjoys a moderate and equable range of temperature throughout the year, although occasionally hot winds may cause the temperature to rise to 100° F. in the eastern part of the State.

(iii) *Comparisons with other Countries.* In respect of Australian temperatures generally, it may be pointed out that the mean annual isotherm for 70° F. extends in South America and South Africa as far south as latitude 33° S., while in Australia it reaches only as far south as latitude 30° S., thus showing that, on the whole, Australia has, latitude for latitude, a more temperate climate than other places in the Southern Hemisphere.

The comparison is even more favourable when the Northern Hemisphere is included, for in the United States of America the 70° F. isotherm extends in several of the western States as far north as latitude 41° N. In Europe, the same isotherm reaches almost to the southern shores of Spain, passing afterwards, however, along the northern shores of Africa till it reaches the Red Sea, when it bends northward along the eastern shore of the Mediterranean till it reaches Syria. In Asia, nearly the whole of the land area south of latitude 40° N. has a mean annual temperature higher than 70° F.

The extreme range of temperature is less than 100° F. over practically the whole of Australia, that figure being only slightly exceeded at a very few places; it is mostly 70° to 90° F. over inland areas, and somewhat less on the coast. In parts of Asia and North America, the extreme range exceeds 130° and 150° F. in some localities.

Along the northern shores of Australia, the temperatures are very equable. At Darwin, for example, the difference in the means for the hottest and coldest month is only 8.4° F., and the extreme readings, or the highest maximum on record and the lowest minimum, show a difference of under 50° F.

The highest temperature recorded in Australia was 127.5° F. at Cloncurry on 16th January, 1889. The world's highest (136° F.) was recorded at Azizia (Tripoli) on 13th August, 1922. The lowest temperature ever recorded in Australia was -8° F. at Charlotte Pass on 14th June, 1945, and again on 22nd July, 1947, as contrasted with the world's lowest recorded temperature of -90° F. at Verkhojansk (Siberia) on 5th and 7th February, 1892.

A comparison of the mean temperatures and the range from the extreme maximum to the extreme minimum temperatures (in whole degrees) of the capital cities of Australia with those of the main cities of some other countries was presented in tabular form in Official Year Book No. 38, page 42.

(iv) *Hottest and Coldest Parts.* A comparison of the temperatures recorded at coast and inland stations shows that, in Australia, as in other continents, the range increases, within certain limits, with increasing distance from the coast.

In the interior of Australia, and during exceptionally dry summers, the temperature occasionally reaches or exceeds 120° F. in the shade. The hottest area of the continent is situated in the northern part of Western Australia about the Marble Bar and Nullagine goldfields, where the maximum shade temperature during the summer sometimes exceeds 100° F. continuously for days and weeks. The longest recorded period was 160 days from 31st October, 1923 to 7th April, 1924.

The area affected and the period of duration of the longest heat waves in Australia are shown in the map and diagram on page 37.

(v) *Tabulated Data for Selected Climatological Stations in Australia.* Tables showing normal mean temperature, extreme temperatures and normal rainfall for each month for

selected climatological stations in each State and the Northern Territory appeared in Official Year Book No. 40, pages 16–23, and similar data for other selected stations in the Commonwealth in Official Year Book No. 39, pages 41–48. Pages 59–66 of this issue contain this information in respect of Canberra, Darwin and the six State capitals.

(vi) *Frosts*.* The Observer's Handbook of the Meteorological Office, London, gives the following definition:—"Injury to the tissues of growing plants is not caused until the temperature has fallen considerably below the freezing point of water (32° F.) and a 'ground frost' is regarded as having occurred when the thermometer on the grass has fallen to 30.4° F. or below".

In Australia, this definition is adopted for stations equipped with terrestrial minimum thermometers. However, these are few in number, so although many rainfall observers record "hoar frost" when seen, for statistical purposes a screen temperature of 36° F. is taken as indicating light frosts at ground level. For heavy frosts, a screen temperature of 32° F. is taken.

In America, a "killing" frost is defined as a frost "that is generally destructive of vegetation". A "black frost" is the phenomenon arising out of a combination of low temperature and low humidity causing rupturing of plant cells by expansion, when freezing takes place, of the water which they contain, though frost crystals are not formed on the ground.

The parts of Australia most subject to low temperature are the eastern highlands from about Omeo in Victoria northward to Cambooya and Bybera in Queensland. Most stations in this region experience more than ten nights per month with readings of 32° F. or under for three to five months of the year. In Tasmania, districts on the Central Plateau are subject to such conditions for three to six months of the year. Minimum temperatures of 32° F. are comparatively infrequent in Western Australia except in parts of the south and south-west. In South Australia, the Yongala district is much more subject to such temperatures than other parts of the State. Much of the south-east of Queensland has a higher frequency of such readings than South Australia. Generally speaking, the frequency is controlled mainly by altitude, latitude and, to a lesser degree, by proximity to the sea.

Frosts may occur within a few miles of the coastline over the whole continent, except in the Northern Territory and a considerable area of Northern Queensland. Regions subject to frost in all months of the year comprise portions of the tabelands of New South Wales, the Eastern Highlands and parts of the Central Divide and Western district in Victoria, practically the whole of Tasmania and a small area in the south-west of Western Australia.

A map showing the average annual number of frost-free days (i.e., days on which the temperature does not fall below 36° F.) appears on page 39.

Over most of the interior of the continent, and on the Highlands in Queensland as far north as the Atherton Plateau frosts appear in April and end in September, but they are infrequent in these months. Minimum temperatures of 32° F. are experienced in most of the sub-tropical interior in June and July.

3. *Humidity*.—After temperature, humidity is the most important element of climate, particularly as regards its effects on human comfort, rainfall supply and conservation and related problems.


"Vapour pressure" is the pressure exerted by the water vapour in the atmosphere. At any given temperature there is a definite upper limit to the amount of water that can exist as vapour in the atmosphere. When this limit is reached, the air is said to be saturated and the pressure of the water vapour is equal to the "saturation vapour pressure".

In this publication, the humidity of the air has been expressed by the relative humidity, which is the quotient of the vapour pressure divided by the saturation vapour pressure and multiplied by one hundred. The mean 9 a.m. relative humidity, as well as its highest and lowest recorded mean values at 9 a.m., are shown in the tables of climatological data for the capital cities (*see* pages 59–66). The mean monthly vapour pressure has also been added to these tables.

The annual curve of vapour pressure derived from the normal monthly values for this element is comparable with the maximum and minimum temperature curves, but the relative humidities, consisting as they do of the extremes for each month, do not show the normal annual fluctuation which would be approximately midway between the extremes.


The order of stations in descending values of 9 a.m. vapour pressure is Darwin, Brisbane Sydney, Perth, Melbourne, Adelaide, Canberra, Hobart and Alice Springs, while the relative humidity at 9 a.m. diminishes in the order, Melbourne, Sydney, Darwin, Brisbane, Canberra, Hobart, Perth, Adelaide and Alice Springs.

* Foley, J. C. Frost in the Australian Region (Bull. 32, 1945).




BUREAU OF METEOROLOGY
**NORMAL DAILY MAXIMUM
 TEMPERATURE**
JANUARY


ISOTHERMS IN DEGREES FAHRENHEIT.




BUREAU OF METEOROLOGY
 NORMAL DAILY MINIMUM
 TEMPERATURE
 JANUARY

ISOTHERMS IN DEGREES FAHRENHEIT






AREA AFFECTED AND PERIOD OF DURATION OF THE LONGEST HEAT WAVES WHEN THE MAXIMUM TEMPERATURE FOR CONSECUTIVE 24 HOURS REACHED OR EXCEEDED 100°F.



NOTE. THE FIGURES ON THE LINES INDICATE THE NUMBER OF CONSECUTIVE DAYS

GREATEST NUMBER OF CONSECUTIVE DAYS ON WHICH THE SHADE TEMPERATURE WAS 100°F OR OVER AT THE PLACES INDICATED





BUREAU OF METEOROLOGY
MAP OF AUSTRALIA
 SHOWING
AVERAGE FROST FREE PERIOD
36° DAYS

SCALE OF MILES
 0 100 200 300

REFERENCE TO DISTRICTS

WESTERN AUSTRALIA		TASMANIA	
1 North Kimberley	21 Perth	1 North	1 Hobart
2 East Kimberley	22 Geraldton	2 Southern	2 Launceston
3 West Kimberley	23 Mandurah	3 Central	3 Devonport
4 Pilbara	24 South Coast	4 North	4 Burnie
5 South Coast	25 South Coast	5 South	5 Launceston
6 North Coast	26 North Coast	6 North	6 Launceston
7 North Coast	27 North Coast	7 South	7 Launceston
8 North Coast	28 North Coast	8 North	8 Launceston
9 North Coast	29 North Coast	9 South	9 Launceston
10 North Coast	30 North Coast	10 North	10 Launceston
11 North Coast	31 North Coast	11 South	11 Launceston
12 North Coast	32 North Coast	12 North	12 Launceston
13 North Coast	33 North Coast	13 South	13 Launceston
14 North Coast	34 North Coast	14 North	14 Launceston
15 North Coast	35 North Coast	15 South	15 Launceston
16 North Coast	36 North Coast	16 North	16 Launceston
17 North Coast	37 North Coast	17 South	17 Launceston
18 North Coast	38 North Coast	18 North	18 Launceston
19 North Coast	39 North Coast	19 South	19 Launceston
20 North Coast	40 North Coast	20 North	20 Launceston
21 North Coast	41 North Coast	21 South	21 Launceston
22 North Coast	42 North Coast	22 North	22 Launceston
23 North Coast	43 North Coast	23 South	23 Launceston
24 North Coast	44 North Coast	24 North	24 Launceston
25 North Coast	45 North Coast	25 South	25 Launceston
26 North Coast	46 North Coast	26 North	26 Launceston
27 North Coast	47 North Coast	27 South	27 Launceston
28 North Coast	48 North Coast	28 North	28 Launceston
29 North Coast	49 North Coast	29 South	29 Launceston
30 North Coast	50 North Coast	30 North	30 Launceston
31 North Coast	51 North Coast	31 South	31 Launceston
32 North Coast	52 North Coast	32 North	32 Launceston
33 North Coast	53 North Coast	33 South	33 Launceston
34 North Coast	54 North Coast	34 North	34 Launceston
35 North Coast	55 North Coast	35 South	35 Launceston
36 North Coast	56 North Coast	36 North	36 Launceston
37 North Coast	57 North Coast	37 South	37 Launceston
38 North Coast	58 North Coast	38 North	38 Launceston
39 North Coast	59 North Coast	39 South	39 Launceston
40 North Coast	60 North Coast	40 North	40 Launceston
41 North Coast	61 North Coast	41 South	41 Launceston
42 North Coast	62 North Coast	42 North	42 Launceston
43 North Coast	63 North Coast	43 South	43 Launceston
44 North Coast	64 North Coast	44 North	44 Launceston
45 North Coast	65 North Coast	45 South	45 Launceston
46 North Coast	66 North Coast	46 North	46 Launceston
47 North Coast	67 North Coast	47 South	47 Launceston
48 North Coast	68 North Coast	48 North	48 Launceston
49 North Coast	69 North Coast	49 South	49 Launceston
50 North Coast	70 North Coast	50 North	50 Launceston
51 North Coast	71 North Coast	51 South	51 Launceston
52 North Coast	72 North Coast	52 North	52 Launceston
53 North Coast	73 North Coast	53 South	53 Launceston
54 North Coast	74 North Coast	54 North	54 Launceston
55 North Coast	75 North Coast	55 South	55 Launceston
56 North Coast	76 North Coast	56 North	56 Launceston
57 North Coast	77 North Coast	57 South	57 Launceston
58 North Coast	78 North Coast	58 North	58 Launceston
59 North Coast	79 North Coast	59 South	59 Launceston
60 North Coast	80 North Coast	60 North	60 Launceston
61 North Coast	81 North Coast	61 South	61 Launceston
62 North Coast	82 North Coast	62 North	62 Launceston
63 North Coast	83 North Coast	63 South	63 Launceston
64 North Coast	84 North Coast	64 North	64 Launceston
65 North Coast	85 North Coast	65 South	65 Launceston
66 North Coast	86 North Coast	66 North	66 Launceston
67 North Coast	87 North Coast	67 South	67 Launceston
68 North Coast	88 North Coast	68 North	68 Launceston
69 North Coast	89 North Coast	69 South	69 Launceston
70 North Coast	90 North Coast	70 North	70 Launceston
71 North Coast	91 North Coast	71 South	71 Launceston
72 North Coast	92 North Coast	72 North	72 Launceston
73 North Coast	93 North Coast	73 South	73 Launceston
74 North Coast	94 North Coast	74 North	74 Launceston
75 North Coast	95 North Coast	75 South	75 Launceston
76 North Coast	96 North Coast	76 North	76 Launceston
77 North Coast	97 North Coast	77 South	77 Launceston
78 North Coast	98 North Coast	78 North	78 Launceston
79 North Coast	99 North Coast	79 South	79 Launceston
80 North Coast	100 North Coast	80 North	80 Launceston





BUREAU OF METEOROLOGY
MAP OF AUSTRALIA
 SHOWING
AVERAGE ANNUAL
EVAPORATION

SCALE OF MILES
 0 100 200 300


FROM EVAPORATION RECORD TO END OF 1946 AND SATURATION DEFICIT RECORDS USING WAITE FORMULA
 $e = 263 S.d.$

(SEE LETTERPRESS ACCOMPANYING)

REFERENCE TO DISTRICTS

WESTERN AUSTRALIA		QUEENSLAND		NEW SOUTH WALES	
1 North Kimberley	30 Central Kimberley	11 Northern Rivers	31 Central Highlands	11 Northern Rivers	31 Central Highlands
2 East Kimberley	32 Central Kimberley	12 North Coast	32 Central Kimberley	12 North Coast	32 Central Kimberley
3 De Grey	33 Upper Western	13 Southern Plains	33 Upper Western	13 Southern Plains	33 Upper Western
4 Fort Pearce	34 Lower Western	14 North West Interior	34 Lower Western	14 North West Interior	34 Lower Western
5 West Coast	35 Pastoral	15 Upper North	35 Pastoral	15 Upper North	35 Pastoral
6 East Coast	36 Pastoral	16 North East	36 Pastoral	16 North East	36 Pastoral
7 North Coast	37 Pastoral	17 Upper North	37 Pastoral	17 Upper North	37 Pastoral
8 Central Coast	38 Pastoral	18 North East	38 Pastoral	18 North East	38 Pastoral
9 South Coast	39 Pastoral	19 Upper North	39 Pastoral	19 Upper North	39 Pastoral
10 Central North	40 Pastoral	20 North East	40 Pastoral	20 North East	40 Pastoral
11 Central South	41 Pastoral	21 Upper North	41 Pastoral	21 Upper North	41 Pastoral
12 South East	42 Pastoral	22 North East	42 Pastoral	22 North East	42 Pastoral
13 North East	43 Pastoral	23 Upper North	43 Pastoral	23 Upper North	43 Pastoral
14 Northern Rivers	44 Pastoral	24 North East	44 Pastoral	24 North East	44 Pastoral
15 Southern Plains	45 Pastoral	25 Upper North	45 Pastoral	25 Upper North	45 Pastoral
16 North West Interior	46 Pastoral	26 North East	46 Pastoral	26 North East	46 Pastoral
17 Upper North	47 Pastoral	27 Upper North	47 Pastoral	27 Upper North	47 Pastoral
18 North East	48 Pastoral	28 North East	48 Pastoral	28 North East	48 Pastoral
19 Upper North	49 Pastoral	29 Upper North	49 Pastoral	29 Upper North	49 Pastoral
20 North East	50 Pastoral	30 North East	50 Pastoral	30 North East	50 Pastoral
21 Upper North	51 Pastoral	31 Central Highlands	51 Pastoral	31 Central Highlands	51 Pastoral
22 North East	52 Pastoral	32 Central Highlands	52 Pastoral	32 Central Highlands	52 Pastoral
23 Upper North	53 Pastoral	33 Upper Western	53 Pastoral	33 Upper Western	53 Pastoral
24 North East	54 Pastoral	34 Lower Western	54 Pastoral	34 Lower Western	54 Pastoral
25 Upper North	55 Pastoral	35 Pastoral	55 Pastoral	35 Pastoral	55 Pastoral
26 North East	56 Pastoral	36 Pastoral	56 Pastoral	36 Pastoral	56 Pastoral
27 Upper North	57 Pastoral	37 Pastoral	57 Pastoral	37 Pastoral	57 Pastoral
28 North East	58 Pastoral	38 Pastoral	58 Pastoral	38 Pastoral	58 Pastoral
29 Upper North	59 Pastoral	39 Pastoral	59 Pastoral	39 Pastoral	59 Pastoral
30 North East	60 Pastoral	40 Pastoral	60 Pastoral	40 Pastoral	60 Pastoral
31 Central Highlands	61 Pastoral	41 Pastoral	61 Pastoral	41 Pastoral	61 Pastoral
32 Central Highlands	62 Pastoral	42 Pastoral	62 Pastoral	42 Pastoral	62 Pastoral
33 Upper Western	63 Pastoral	43 Pastoral	63 Pastoral	43 Pastoral	63 Pastoral
34 Lower Western	64 Pastoral	44 Pastoral	64 Pastoral	44 Pastoral	64 Pastoral
35 Pastoral	65 Pastoral	45 Pastoral	65 Pastoral	45 Pastoral	65 Pastoral
36 Pastoral	66 Pastoral	46 Pastoral	66 Pastoral	46 Pastoral	66 Pastoral
37 Pastoral	67 Pastoral	47 Pastoral	67 Pastoral	47 Pastoral	67 Pastoral
38 Pastoral	68 Pastoral	48 Pastoral	68 Pastoral	48 Pastoral	68 Pastoral
39 Pastoral	69 Pastoral	49 Pastoral	69 Pastoral	49 Pastoral	69 Pastoral
40 Pastoral	70 Pastoral	50 Pastoral	70 Pastoral	50 Pastoral	70 Pastoral
41 Pastoral	71 Pastoral	51 Pastoral	71 Pastoral	51 Pastoral	71 Pastoral
42 Pastoral	72 Pastoral	52 Pastoral	72 Pastoral	52 Pastoral	72 Pastoral
43 Pastoral	73 Pastoral	53 Pastoral	73 Pastoral	53 Pastoral	73 Pastoral
44 Pastoral	74 Pastoral	54 Pastoral	74 Pastoral	54 Pastoral	74 Pastoral
45 Pastoral	75 Pastoral	55 Pastoral	75 Pastoral	55 Pastoral	75 Pastoral
46 Pastoral	76 Pastoral	56 Pastoral	76 Pastoral	56 Pastoral	76 Pastoral
47 Pastoral	77 Pastoral	57 Pastoral	77 Pastoral	57 Pastoral	77 Pastoral
48 Pastoral	78 Pastoral	58 Pastoral	78 Pastoral	58 Pastoral	78 Pastoral
49 Pastoral	79 Pastoral	59 Pastoral	79 Pastoral	59 Pastoral	79 Pastoral
50 Pastoral	80 Pastoral	60 Pastoral	80 Pastoral	60 Pastoral	80 Pastoral
51 Pastoral	81 Pastoral	61 Pastoral	81 Pastoral	61 Pastoral	81 Pastoral
52 Pastoral	82 Pastoral	62 Pastoral	82 Pastoral	62 Pastoral	82 Pastoral
53 Pastoral	83 Pastoral	63 Pastoral	83 Pastoral	63 Pastoral	83 Pastoral
54 Pastoral	84 Pastoral	64 Pastoral	84 Pastoral	64 Pastoral	84 Pastoral
55 Pastoral	85 Pastoral	65 Pastoral	85 Pastoral	65 Pastoral	85 Pastoral
56 Pastoral	86 Pastoral	66 Pastoral	86 Pastoral	66 Pastoral	86 Pastoral
57 Pastoral	87 Pastoral	67 Pastoral	87 Pastoral	67 Pastoral	87 Pastoral
58 Pastoral	88 Pastoral	68 Pastoral	88 Pastoral	68 Pastoral	88 Pastoral
59 Pastoral	89 Pastoral	69 Pastoral	89 Pastoral	69 Pastoral	89 Pastoral
60 Pastoral	90 Pastoral	70 Pastoral	90 Pastoral	70 Pastoral	90 Pastoral
61 Pastoral	91 Pastoral	71 Pastoral	91 Pastoral	71 Pastoral	91 Pastoral
62 Pastoral	92 Pastoral	72 Pastoral	92 Pastoral	72 Pastoral	92 Pastoral
63 Pastoral	93 Pastoral	73 Pastoral	93 Pastoral	73 Pastoral	93 Pastoral
64 Pastoral	94 Pastoral	74 Pastoral	94 Pastoral	74 Pastoral	94 Pastoral
65 Pastoral	95 Pastoral	75 Pastoral	95 Pastoral	75 Pastoral	95 Pastoral
66 Pastoral	96 Pastoral	76 Pastoral	96 Pastoral	76 Pastoral	96 Pastoral
67 Pastoral	97 Pastoral	77 Pastoral	97 Pastoral	77 Pastoral	97 Pastoral
68 Pastoral	98 Pastoral	78 Pastoral	98 Pastoral	78 Pastoral	98 Pastoral
69 Pastoral	99 Pastoral	79 Pastoral	99 Pastoral	79 Pastoral	99 Pastoral
70 Pastoral	100 Pastoral	80 Pastoral	100 Pastoral	80 Pastoral	100 Pastoral
71 Pastoral	101 Pastoral	81 Pastoral	101 Pastoral	81 Pastoral	101 Pastoral
72 Pastoral	102 Pastoral	82 Pastoral	102 Pastoral	82 Pastoral	102 Pastoral
73 Pastoral	103 Pastoral	83 Pastoral	103 Pastoral	83 Pastoral	103 Pastoral
74 Pastoral	104 Pastoral	84 Pastoral	104 Pastoral	84 Pastoral	104 Pastoral
75 Pastoral	105 Pastoral	85 Pastoral	105 Pastoral	85 Pastoral	105 Pastoral
76 Pastoral	106 Pastoral	86 Pastoral	106 Pastoral	86 Pastoral	106 Pastoral
77 Pastoral	107 Pastoral	87 Pastoral	107 Pastoral	87 Pastoral	107 Pastoral
78 Pastoral	108 Pastoral	88 Pastoral	108 Pastoral	88 Pastoral	108 Pastoral
79 Pastoral	109 Pastoral	89 Pastoral	109 Pastoral	89 Pastoral	109 Pastoral
80 Pastoral	110 Pastoral	90 Pastoral	110 Pastoral	90 Pastoral	110 Pastoral



PROJECTION - SIMPLE CONIC. STANDARD PARALLEL - 25 S.



COMMONWEALTH OF AUSTRALIA.
 BUREAU OF METEOROLOGY MELBOURNE.
**MONTHLY DISTRIBUTION OF RAINFALL
 OVER AUSTRALIA.**
 REVISED TO END OF 1939.

SCALE OF MILES
 100 50 0 100 200 300

EXPLANATION— THE GRAPHS SHOW THE AVERAGE RAINFALL IN EACH MONTH OF THE YEAR IN THE VARIOUS RAINFALL DISTRICTS, THE BOUNDARIES OF WHICH ARE SHOWN BY THE BROKEN LINES ON THE MAP. THE NUMBERS ON THE MAP REFER TO THE DISTRICTS. THE CORRESPONDING DISTRICT NAMES ARE GIVEN IN THE INSERT ON PAGE 49.
 THE COLUMNS FOR THE SUCCESSIVE MONTHS, JANUARY TO DECEMBER, RUN FROM LEFT TO RIGHT.
 THE LENGTH OF A VERTICAL COLUMN REPRESENTS THE AVERAGE MONTHLY RAINFALL ACCORDING TO THE SCALE GIVEN BELOW.




ISSUED UNDER THE AUTHORITY OF THE MINISTER OF STATE FOR THE DEPARTMENT OF AIR
BY H.N. WARREN DIRECTOR OF METEOROLOGICAL SERVICES

9 A.M.

LOCAL MEAN TIME

WIND ROSES

SHOWING MONTHLY FREQUENCY
DISTRIBUTION OF WIND DIRECTIONS
TO EIGHT POINTS OF THE COMPASS




ISSUED UNDER THE AUTHORITY OF THE MINISTER OF STATE FOR THE DEPARTMENT OF AIR
BY H.N. WARREN DIRECTOR OF METEOROLOGICAL SERVICES

3 P.M.

LOCAL MEAN TIME

WIND ROSES

SHOWING MONTHLY FREQUENCY
DISTRIBUTION OF WIND DIRECTIONS
TO EIGHT POINTS OF THE COMPASS



Further reference to humidity will be found in the section on effective temperature (see p. 29).

4. **Evaporation.**—(i) *General.* The rate and quantity of evaporation in any territory is influenced by the prevailing temperature, and by atmospheric humidity, pressure and wind movement. In Australia, the question is, perhaps, of more than ordinary importance, since in its drier regions water has often to be conserved in tanks and dams. The magnitude of the economic loss by evaporation will be appreciated from the map reproduced herein (see p. 40) which shows that the yearly amount varies from about 20 inches over western Tasmania to more than 100 inches over the central and north-western parts of Australia. Over an area of 70 per cent. of the continent, comprising most inland districts and extending to the coast in the North-West and Eucla divisions of Western Australia, the rainfall does not exceed the evaporation during any month of the year. The central and north-western portions of the continent, comprising 46 per cent. of the total land mass, experience evaporation more than twice as great as their rainfall; it is noteworthy that the vegetation over most of this region is characterized by acacia, semi-desert, shrub steppe and porcupine grass. Since the loss by evaporation depends largely on the exposed area, tanks and dams so designed that the surface shall be a minimum are advantageous. Further, the more they are protected from the direct rays of the sun and from winds by means of suitable tree planting, the less will be the loss by evaporation. The Mansfield Process for treatment of tanks, dams and ponds by hexadecanol film, materially reducing effective evaporation, is a recent development which is already having beneficial results. These matters are naturally of more than ordinary concern in the drier districts of Australia.

(ii) *Comments on Map of Average Annual Evaporation.* The map of average annual evaporation in Australia (see p. 40) has been compiled on the basis of records obtained from a number of evaporimeters supplemented by estimates derived from records of saturation deficit by applying the Waite Institute factor of 263.* Some modification of the latter values was found to be necessary in comparison with recordings of evaporimeters.

The standard evaporation tank used in Australia is cylindrical in form and is 36 inches in diameter and 36 inches deep. It is surrounded by a 6-inch water jacket and the whole is sunk into the ground so that the water surface is approximately at ground level.

Saturation deficit is obtained from readings of dry and wet bulb thermometers exposed in a standard Stevenson thermometer shelter. Saturation deficit is the difference between the vapour pressure indicated by the dry and wet bulb readings, and the saturation vapour pressure corresponding to the dry bulb temperature.

The Waite formula, $e = 263 \text{ s.d.}$, is not an exact relationship, but it takes account of one of the major factors in evaporation, i.e., the difference between saturation vapour pressures at the mean dew point and at the mean air temperature. Errors in the formula are found to be fairly consistent in considerable areas of Australia and corrections have been applied accordingly. No evaporation records are available north of latitude 20° , and corrections have been extrapolated for these areas. The evaporation stations on which estimates for the tropics have been based are Alice Springs (Northern Territory) and Winton (Queensland), and to a lesser degree Blackall (Queensland) and Marble Bar (Western Australia).

The map thus presents an estimate of evaporation for which allowance should be made for a certain margin of error (perhaps 10 per cent. or so) on the conservative side. In the absence of definite information, such a map should serve a useful purpose as a basis for many climatic studies.

For graphs and tables of mean monthly evaporation and rainfall at certain selected stations, see Official Year Book No. 37, pages 34–35.

5. **Rainfall.**—(i) *General.* The rainfall of any region is determined mainly by the direction and route of the prevailing winds, by the varying temperatures of the earth's surface over which they blow, and by its physiographical features.

Australia lies within the zones of the south-east trades and "prevailing" westerly winds. The southern limit of the south-east trades strikes the eastern shores at about 30° south latitude, and, with very few exceptions, the heaviest rains of the Australian continent are precipitated along the Pacific slopes to the north of that latitude, the varying quantities being more or less regulated by the differences in elevation of the shores and of the chain of mountains from the New South Wales northern border to Thursday Island,

* Prescott, J. A. "Atmospheric Saturation Deficit in Australia" (Trans. Royal Society, S.A., Vol. LV., 1931).

upon which the rain-laden winds blow. The converse effect is exemplified on the north-west coast of Western Australia, where the prevailing winds, blowing from the interior of the continent instead of from the ocean, result in the lightest coastal rain in Australia.

The westerly winds, which skirt the southern shores, are responsible for the reliable, generally light to moderate rains enjoyed by the south-western portion of Western Australia, the agricultural areas of South Australia, a great part of Victoria, and the whole of Tasmania.

(ii) *Distribution of Rainfall.* The average annual rainfall map of Australia (*see* p. 41) shows that the heaviest yearly falls occur on the north coast of Queensland (up to more than 160 inches) and in western Tasmania (up to 140 inches), while from 50 to over 60 inches are received on parts of the eastern seaboard from Jervis Bay to the northern part of Cape York Peninsula, also around Darwin, on the West Kimberley coast, near Cape Leeuwin, about the Australian Alps in eastern Victoria and New South Wales, and on the north-eastern highlands in Tasmania. A great part of the interior of the continent, stretching from the far west of New South Wales and the south-west of Queensland to the vicinity of Shark Bay in Western Australia, has a very low average rainfall of less than 10 inches a year. Between these two regions of heavy and very low rainfall are the extensive areas which experience useful to good rains, and in the southern and eastern parts of which are found the best country and most of the population and primary production.

(iii) *Factors Determining Occurrence, Intensity and Seasonal Distribution of Rainfall.* Reference has already been made to the frequent rains occurring in the north-eastern coastal districts of Queensland with the prevailing south-east trade winds and to similar rains in the west of Tasmania with the prevailing westerly winds. Other rains in Australia are associated mainly with tropical and southern depressions.

The former chiefly affect the northern, eastern, and to some extent the central parts of the continent and operate in an irregular manner during the warmer half of the year, but principally from December to March. They vary considerably in activity and scope from year to year, occasionally developing into severe storms off the east and north-west coasts. Tropical rainstorms sometimes cover an extensive area, half of the continent on occasions receiving moderate to very heavy falls during a period of a few days. Rain is also experienced, with some regularity, with thunderstorms in tropical areas, especially near the coast. All these tropical rains, however, favour mostly the northern and eastern parts of the area referred to; the other parts further inland receive lighter, less frequent and less reliable rainfall. With the exception of districts near the east coast, where some rain falls in all seasons, the tropical parts of the continent receive useful rains only on rare occasions from May to September.

The southern depressions are most active in the winter—June to August—and early spring months. The rains associated with them are fairly reliable and frequent over southern Australia and Tasmania, and provide during that period the principal factor in the successful growing of wheat. These depressions also operate with varying activity during the remainder of the year, but the accompanying rains are usually lighter. The southern rains favour chiefly the south-west of Western Australia, the agricultural districts of South Australia, Victoria, Tasmania, and the southern parts of New South Wales. They sometimes extend into the drier regions of the interior, but only infrequently and irregularly.

The map showing mean monthly distribution of rainfall over Australia (*see* p. 42) gives, in graphic form, information on the amount and occurrence of rain.

(iv) *Wettest and Driest Regions.* The wettest known part of Australia is on the north-east coast of Queensland, between Port Douglas and Cardwell, where Deeral on the northern coast-line has an average annual rainfall of 172.26 inches and Tully on the Tully River 179.26 inches. In addition, three stations situated on, or adjacent to, the Johnstone and Russell Rivers have an average annual rainfall of between 144 and 169 inches. The maximum and minimum annual amounts there, in inches, are:—Deeral, 287.18 in 1945 and 94.65 in 1951, a range of 192.53 inches; Tully, 310.92 in 1950 and 104.98 in 1943, a range of 205.94 inches; Goondi, 241.53 in 1894 and 67.88 in 1915, a range of 173.65 inches; Innisfail, 232.06 in 1950 and 69.87 in 1902, a range of 162.19 inches; Harvey Creek, 254.77 in 1921 and 80.47 in 1902, a range of 174.30 inches.

On five occasions, more than 200 inches have been recorded at Goondi, the last of these being in 1950, when 204.97 inches were registered. The records at this station cover a period of 70 years.

In 30 years of record, Tully has exceeded 200 inches on eleven occasions, and in 28 complete years of record Harvey Creek has exceeded this figure four times.

In Tasmania, the wettest part is in the West Coast region, the average annual rainfall at Lake Margaret being 146.51 inches, with a maximum of 177.30 inches in 1948.

The driest known part of the continent is in an area of approximately 180,000 square miles surrounding Lake Eyre in South Australia, where the annual average is between 4 and 6 inches and where the fall rarely exceeds 10 inches in twelve months.

Records at stations have at times been interrupted, but of the 23 stations in this region which have an annual average of less than 5 inches, six have complete records extending from 30 to 55 years. Of these, Mulka has the lowest average of 4.05 inches (34 years), followed by Troudaninna with an average of 4.15 inches over 42 years. Troudaninna in the period 1893 to 1936 had only one year in which the total exceeded 9 inches (11.07 inches in 1894). There have been protracted periods when the average has been even less than 3 inches. From 1895 to 1903, Troudaninna received the following annual totals:— 2.78, 0.99, 5.71, 3.04, 3.18, 2.83, 1.80, 1.11, 4.87, an average of 2.91 inches. From 1918 to 1929 the average was only 2.65 inches, and in this period from December, 1924, to November, 1929, the average was only 1.70 inches.

Mulka since 1918 has only twice exceeded 10 inches for the annual total (11.72 inches in 1920 and 13.56 in 1955) and on 16 occasions in 34 years the annual total has been less than 3 inches. In one particular period from October, 1926 to September, 1930, the average was only 1.26 inches (505 points in 48 months). However, at Kanowana, an even lower four-year average of 1.12 inches was recorded between 1896 and 1899 with yearly totals of 43, 225, 87 and 94 points. An even smaller yearly total was recorded at Mungerina in 1889 when only 39 points were recorded on five days.

The average number of days of rain per month in this region is only 1 to 2 and the annual number ranges between 10 and 20. Oodnadatta (standard 30 years' average rainfall equal to 4.44 inches) has an average of 20 days of rain per year, while Cordillo Downs in the extreme north-east corner of the State of South Australia receives 5.16 inches on twelve days per year, averaging about one day of rain each month in the 30 year period 1911-1940.

No part of the earth, so far as is known, is absolutely rainless, and although at Arica, in northern Chile, the rainfall over a period of 15 years was nil, a further two years in which there were three measurable showers made the "average" for 17 years 0.02 inches.

(v) *Quantities and Distribution of Rainfall.* The general distribution is best seen from the rainfall map (see page 41) which shows the areas subject to average annual rainfalls lying between certain limits. The proportions of the total area of each State and of Australia as a whole enjoying varying quantities of rainfall determined from the latest available information are shown in the following table:—

As per 1957. **AVERAGE ANNUAL RAINFALL DISTRIBUTION.**
(Per Cent.)

Average Annual Rainfall.	W. Aust.	N. Terr.	S. Aust.	Q'land.	N.S.W. (a)	Vic.	Tas. (b)	Total.
Under 10 inches ..	58.0	24.7	82.8	13.0	19.7	Nil	Nil	37.6
10 and under 15 ins.	22.4	32.4	9.4	14.4	23.5	22.4	Nil	19.9
15 and under 20 ..	6.8	9.7	4.5	19.7	17.5	15.2	0.7	10.9
20 and under 25 ..	3.7	6.6	2.2	18.8	14.2	17.9	11.0	9.1
25 and under 30 ..	3.7	9.3	0.8	11.6	9.1	18.0	11.4	7.3
30 and under 40 ..	3.3	4.7	0.3	11.1	9.9	16.1	20.4	6.6
40 inches and over ..	2.1	12.6	Nil	11.4	6.1	10.4	56.5	8.6
Total ..	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

(a) Includes Australian Capital Territory.
are available.

(b) Over an area of 2,777 square miles no records

Referring first to the capital cities, the records of which are given in the next table, it will be seen that Sydney, with an average rainfall of 44.80 inches, occupies the chief place; Brisbane, Perth, Melbourne, Hobart and Canberra follow in that order, Adelaide with 21.09 inches being the driest. The extreme range from the wettest to the driest year is greatest at Brisbane (72.09 inches) and least at Adelaide (19.46 inches).

In order to show how the rainfall is distributed throughout the year in various parts of the continent, average figures for the various climatological districts have been selected (see map on page 42). The figures for Northern Rivers (District 14), show that nearly

the whole of the rainfall occurs there in the summer months, while little or none falls in the middle of the year. The figures for the Central Coast, south-west of Western Australia (District 9), are the reverse, for while the summer months are dry, the winter months are very wet. In the districts containing Melbourne and Hobart, the rain is fairly well distributed throughout the twelve months, with a maximum in October for both districts. In Queensland, the heaviest rains fall in the summer months, but good averages are also maintained during the other seasons in eastern parts.

On the coast of New South Wales, the first half of the year is the wettest, with heaviest falls in the autumn; the averages during the last six months are fair, and moderately uniform. Generally, it may be said that approximately one-third of the area of the continent, principally in the eastern and northern parts, enjoys an annual average rainfall of from 20 to 50 inches or more, the remaining two-thirds averaging from 5 to 20 inches.

(vi) *Tables of Rainfall.* The following table of rainfall for a fairly long period of years for each of the Australian capitals affords information as to the variability of the fall in successive years, and the list which follows in the next paragraph of the more remarkable falls furnishes information as to what may be expected on particular occasions.

RAINFALL: AUSTRALIAN CAPITAL CITIES.

Year.	PERTH.		ADELAIDE.		BRISBANE.		SYDNEY.		CANNBERRA.(a)		MELBOURNE.		HOBART.(b)	
	Amount.	No. of Days.	Amount.	No. of Days.	Amount.	No. of Days.	Amount.	No. of Days.	Amount.	No. of Days.	Amount.	No. of Days.	Amount.	No. of Days.
1927 ..	36.59	133	16.92	101	62.08	130	48.56	138	18.59	90	17.98	135	20.13	185
1928 ..	44.88	140	19.43	107	52.64	145	40.07	130	23.12	70	24.09	151	30.23	205
1929 ..	36.77	132	17.51	119	39.78	118	57.90	129	28.81	90	28.01	168	26.55	194
1930 ..	39.80	129	18.65	116	41.22	144	44.47	141	17.33	82	25.41	145	19.38	152
1931 ..	39.18	118	22.26	145	66.72	136	49.22	153	24.02	103	28.63	164	27.17	179
1932 ..	39.40	121	25.04	141	24.79	97	37.47	146	20.18	118	31.08	179	30.29	155
1933 ..	32.47	116	22.12	130	49.71	118	42.71	153	20.78	96	22.28	136	23.18	182
1934 ..	40.61	120	20.24	125	54.26	117	64.91	183	35.58	131	33.53	157	23.17	194
1935 ..	32.28	129	23.45	140	34.64	111	30.97	131	23.78	95	29.98	183	32.22	196
1936 ..	30.64	118	19.34	121	21.77	101	30.22	130	26.24	108	24.30	187	19.60	178
1937 ..	35.28	120	23.01	128	34.79	113	52.00	157	20.46	82	21.45	144	20.65	160
1938 ..	29.64	111	19.26	119	43.49	110	39.17	132	19.26	79	17.63	131	31.32	169
1939 ..	45.70	123	23.29	139	41.43	122	33.67	127	27.63	116	33.11	166	27.23	188
1940 ..	20.00	98	16.16	116	42.37	93	39.34	125	17.38	64	19.83	126	17.17	135
1941 ..	34.74	122	22.56	126	31.50	105	26.74	129	19.55	91	31.78	157	23.49	145
1942 ..	39.24	140	25.44	133	44.01	125	48.29	121	25.76	104	29.79	148	19.42	163
1943 ..	31.46	117	17.84	135	50.68	126	50.74	136	24.59	123	18.80	150	20.84	149
1944 ..	27.39	123	17.13	114	27.85	100	31.04	115	12.05	75	21.32	143	26.23	151
1945 ..	52.67	137	17.85	105	48.16	130	46.47	136	22.35	100	19.22	152	16.92	157
1946 ..	41.47	122	22.59	135	38.66	83	36.05	111	22.31	94	29.80	177	39.45	193
1947 ..	43.42	137	21.89	146	60.30	146	41.45	137	27.95	135	30.47	163	38.61	181
1948 ..	34.75	126	21.40	122	41.54	106	38.83	131	32.11	101	20.98	155	23.42	178
1949 ..	27.15	126	18.23	119	47.18	121	66.26	149	27.71	100	31.41	163	22.85	157
1950 ..	32.27	122	16.06	91	63.93	152	86.33	183	43.35	132	26.18	147	19.25	131
1951 ..	34.14	127	25.44	135	33.89	87	53.15	143	22.00	103	29.85	155	24.57	163
1952 ..	39.28	123	19.99	128	33.49	122	59.19	130	37.87	141	34.39	177	30.35	165
1953 ..	37.14	119	20.00	121	43.60	101	40.86	110	19.40	102	28.38	148	28.06	162
1954 ..	28.05	112	16.73	109	61.36	142	41.29	134	18.81	82	33.53	139	27.20	143
1955 ..	46.52	138	24.58	134	50.41	136	72.46	160	30.85	124	30.70	160	22.32	168
1956 ..	37.35	107	27.24	154	59.18	120	67.33	155	40.46	150	30.96	188	36.63	175
1957 ..	33.40	117	16.71	110	20.58	80	27.13	110	14.41	81	20.68	146	28.66	129
Average	35.05	121	21.04	121	44.82	124	47.20	150	24.53	103	25.90	143	24.91	167
No. of														
Stand-	82	82	119	119	106	98	99	99	30	30	102	102	75	75
ard 30														
years'														
Normal(c)	35.99	128	21.09	122	40.09	117	44.80	143			25.89	156	25.03	180

(a) Commonwealth Forestry Bureau; records in issues of the Official Year Book prior to No. 36 were for the station at Acton which closed down in 1939. (b) Records taken from present site commenced 1883. (c) 1911-1940.

6. Remarkable Falls of Rain.—The following are the most notable falls of rain which have occurred within a period of twenty-four hours ending at 9 a.m. in the various States and Territories. For other very heavy falls at various localities, reference may be made to Official Year Book No. 14, pages 60–64, No. 22, pages 46–48 and No. 29, pages 43, 44 and 51.

HEAVY RAINFALLS: WESTERN AUSTRALIA, UP TO 1957 INCLUSIVE.

Name of Town or Locality.	Date.	Amt.	Name of Town or Locality.	Date.	Amt.
		in.			in.
Whim Creek ..	3 Apr., 1898	29.41	Pilbara ..	2 Apr., 1898	14.04
Fortesque ..	3 May, 1890	23.36	Roebuck Plains ..	5 Jan., 1917	14.01
Roebuck Plains ..	6 Jan., 1917	22.36	Broome ..	6 Jan., 1917	14.00
Widjip ..	1 Apr., 1934	19.54	Carlton Hill ..	7 Feb., 1942	12.75
Derby ..	7 Jan., 1917	16.47	Towrana ..	1 Mar., 1943	12.16
Boodarie ..	21 Mar., 1899	14.53	Marble Bar ..	2 Mar., 1941	12.00
Balla Balla ..	21 Mar., 1899	14.40	Jimba Jimba ..	1 Mar., 1943	11.54
Winderrie ..	17 Jan., 1923	14.23			

HEAVY RAINFALLS: NORTHERN TERRITORY, UP TO 1957, INCLUSIVE.

Name of Town or Locality.	Date.	Amt.	Name of Town or Locality.	Date.	Amt.
		in.			in.
Brocks Creek ..	24 Dec., 1915	14.33	Cape Don ..	13 Jan., 1935	13.58
Groote Eylandt ..	9 Apr., 1931	14.29	Bathurst Island		
Borroloola ..	14 Mar., 1899	14.00	Mission ..	7 Apr., 1925	11.85
Timber Creek ..	5 Feb., 1942	13.65	Darwin ..	7 Jan., 1897	11.67

HEAVY RAINFALLS: SOUTH AUSTRALIA, UP TO 1957, INCLUSIVE.

Name of Town or Locality.	Date.	Amt.	Name of Town or Locality.	Date.	Amt.
		in.			in.
Ardrrossan ..	18 Feb., 1946	8.10	Wilmington ..	1 Mar., 1921	7.12
Carpa ..	18 Feb., 1946	7.83	Port Victoria ..	18 Feb., 1946	7.08
Wynbring ..	28 Feb., 1921	7.70	Mannum ..	25 Jan., 1941	6.84
Edithburg ..	18 Feb., 1946	7.46	Wirrabarra ..	7 Mar., 1910	6.80
Hesso ..	18 Feb., 1946	7.36	Cape Willoughby	18 Feb., 1946	6.80
Maitland ..	18 Feb., 1946	7.21	Torrens Vale ..	25 Jan., 1941	6.77

HEAVY RAINFALLS: QUEENSLAND, UP TO 1957, INCLUSIVE.

Name of Town or Locality.	Date.	Amt.	Name of Town or Locality.	Date.	Amt.
		in.			in.
Crohamhurst ..	3 Feb., 1893	35.71	Flat Top Island ..	21 Jan., 1918	25.18
Port Douglas ..	1 Apr., 1911	31.53	Landsborough ..	3 Feb., 1893	25.15
Yarrabah ..	2 Apr., 1911	30.65	Harvey Creek ..	31 Jan., 1913	24.72
Mooloolah ..	3 Feb., 1893	29.11	Kuranda ..	1 Apr., 1911	24.30
Kuranda ..	2 Apr., 1911	28.80	Babinda (Cairns)	2 Mar., 1935	24.14
Harvey Creek ..	3 Jan., 1911	27.75	Goondi ..	30 Jan., 1913	24.10
Sarina ..	26 Feb., 1913	27.75	Banyan (Cardwell)	12 Feb., 1927	24.00
Plane Ck. (Mackay)	26 Feb., 1913	27.73	Carruchan ..	24 Jan., 1934	24.00
Deeral ..	2 Mar., 1935	27.60	Tully Mill ..	12 Feb., 1927	23.86
Yarrabah Mission	24 Jan., 1916	27.20	Macnade Mill ..	7 Jan., 1901	23.33
Springbrook ..	24 Jan., 1947	27.07	Woodlands (Yepoon)		
Springbrook ..	21 Feb., 1954	27.04		1 Feb., 1893	23.07
Buderim Mountain	12 Jan., 1898	26.20			

HEAVY RAINFALLS: NEW SOUTH WALES, UP TO 1957, INCLUSIVE.

Name of Town or Locality.	Date.	Amt.	Name of Town or Locality.	Date.	Amt.
		in.			in.
Dorrigo ..	24 June, 1950	25.04	Viaduct Creek ..	15 Mar., 1936	20.00
Cordeaux River ..	14 Feb., 1898	22.58	Buladelah ..	16 Apr., 1927	19.80
Morpeth ..	9 Mar., 1893	21.52	Orara Upper ..	24 June, 1950	19.80
Broger's Creek ..	13 Jan., 1911	20.83	Madden's Creek ..	13 Jan., 1911	18.68
South Head (Sydney Harbour) ..	16 Oct., 1844	20.41	Condong ..	27 Mar., 1887	18.66
" "	29 Apr., 1841	20.12	Candelo ..	27 Feb., 1919	18.58
Mount Pleasant ..	5 May, 1925	20.10	Mt. Kembla ..	13 Jan., 1911	18.25
Broger's Creek ..	14 Feb., 1898	20.05	Bega ..	27 Feb., 1919	17.88
Towamba ..	5 Mar., 1893	20.00	Kembla Heights	13 Jan., 1911	17.46
			Foxground ..	11 Sept., 1950	17.04

HEAVY RAINFALL: AUSTRALIAN CAPITAL TERRITORY, UP TO 1957, INCLUSIVE.

Name of Town or Locality.	Date.	Amt.	Name of Town or Locality.	Date.	Amt.
		in.			in.
Cotter Junction ..	27 May, 1925	7.13	Uriarra (Woodside)	27 May, 1925	6.57
Canberra (Acton)	27 May, 1925	6.84	Land's End ..	27 May, 1925	6.35

HEAVY RAINFALLS: VICTORIA, UP TO 1957, INCLUSIVE.

Name of Town or Locality.	Date.	Amt.	Name of Town or Locality.	Date.	Amt.
		in.			in.
Balook ..	18 Feb., 1951	10.81	Blackwood (Greenhill) ..	26 Jan., 1941	8.98
Hazel Park ..	1 Dec., 1934	10.50	Tambo Crossing ..	13 July, 1925	8.89
Kalorama ..	1 Dec., 1934	10.05	Corinella ..	28 June, 1948	8.75
Cann River ..	16 Mar., 1938	9.94	Erica ..	1 Dec., 1934	8.66
Tonghi Creek ..	27 Feb., 1919	9.90	Mt. Buffalo ..	6 June, 1917	8.53
Cann River ..	27 Feb., 1919	9.56	Korumburra ..	1 Dec., 1934	8.51
Olinda ..	1 Dec., 1934	9.10			

HEAVY RAINFALLS: TASMANIA, UP TO 1957, INCLUSIVE.

Name of Town or Locality.	Date.	Amt.	Name of Town or Locality.	Date.	Amt.
		in.			in.
Mathinna ..	5 Apr., 1929	13.25	Riana ..	5 Apr., 1929	11.08
Cullenswood ..	5 Apr., 1929	11.12	Triabunna ..	5 June, 1923	10.20

7. **Snowfall.**—Light snow has been known to fall occasionally as far north as latitude 31° S., and from the western to the eastern shores of the continent. During exceptional seasons, it has fallen simultaneously over two-thirds of New South Wales, and has extended at times along the whole of the Great Dividing Range, from its southern extremity in Victoria as far north as Toowoomba in Queensland. During the winter, for several months, snow covers the ground to a great extent on the Australian Alps, where the temperature falls below zero Fahrenheit during the night. In the ravines around Mt. Kosciusko and similar localities the snow never entirely disappears after a severe winter.

8. **Hail.**—Hail falls most frequently along the southern shores of the continent in the winter, and over eastern Australia during the summer months. The size of the hailstones generally increases with distance from the coast. A summer rarely passes

without some station experiencing a fall of stones exceeding in size an ordinary hen egg and many riddled sheets of light-gauge galvanized iron bear evidence of the weight and penetrating power of the stones.

The hailstones occur most frequently when the barometric readings indicate a flat and unstable condition of pressure. Tornadoes or tornadic tendencies are almost invariably accompanied by hail, and on the east coast the clouds from which the stones fall are frequently of a remarkable sepia-coloured tint.

9. Droughts.—(a) *General*. The following extracts are taken from a Bureau of Meteorology publication *Droughts in Australia* by J. C. Foley (Bulletin 43, 1957). Sections of this publication on the definition of drought, drought severity according to climate, and general descriptions of major widespread Australian droughts are given fully but the method of treatment of data and assessment of the severity of droughts is given in outline only. Droughts which were confined to comparatively small areas of Australia are not mentioned below but are treated in detail in Bulletin 43.

(b) *The Definition of Drought*. A commonly accepted broad definition is "dryness due to lack of rain". Various writers have used this definition but for purposes of comparison a more precise definition is required. For instance, one writer represents drought frequencies by the number of consecutive days without rain experienced in various periods of years. Others prefer a definition in terms of deviation from the normal rainfall e.g. a defect of 50% from the mean rainfall of three or more consecutive months. Again, more recent writers have preferred to consider drought in terms of the effect of rainfall deficiencies on the plant life of an area, especially upon plants of economic importance. The views of Russell and Blair are accepted as a basis for this study. Russell, a pioneer meteorologist (Government Astronomer) in New South Wales, based his definition on his experiences of the droughts of 1888 and 1895. He says "The word drought is not used here" (in Australia) "as in the sense in which it is often used in England and elsewhere, that is, signifying a period of a few days or weeks in which not a drop of rain falls, but it is used to signify a period of months or years during which little rain falls and the country gets burnt up, grass and water disappear, crops become worthless and sheep and cattle die". Blair regards drought as a relative term. As crops in different parts of the world are more or less adjusted to the normal moisture conditions and to the normal rainfall distribution for the region and a rainless period that would constitute a drought in one region may not be unusual or injurious in another. To these views it may be added that crops and other vegetation tolerate variations in moisture conditions within certain limits and a drought occurs when the lower limit of moisture which can be tolerated is exceeded.

It is therefore considered sufficient and justifiable to consider a drought to have occurred when there has been a persistent deficiency of rainfall over a period of months without specifying precisely the degree of deficiency or its duration. Of course, the effects of rainfall deficiency depend on other factors such as the moisture status of the ground and the state of water supplies when a period of deficiency sets in, the season of the year, temperature, evaporation and wind conditions and the resistance to drought of various types of crops and other vegetation. However it is not practicable or necessary to take these factors into account in a study which aims to discuss periods of serious rainfall deficiency in the meteorological history of Australia and their effects on primary production as shown by the reports available.

(c) *The Treatment of Data and Assessment of Degree of Severity*. The problem was approached primarily from the viewpoint of deficiency of rainfall compared with the average for all years of record. Periods of rainfall deficiency were identified by means of a "residual mass" or "cumulated residual" graph, which is a continuous plot of cumulative totals of departure from the average, plus and minus, of monthly rainfall through all years of record. Residual mass graphs show clearly the beginning and end of periods of rainfall deficiency and excess and the amount of deficiency in drought periods. For construction of the residual mass graphs, 28 areas of agricultural or pastoral importance were selected, with a group of five or ten stations with a long series of records to represent the rainfall regime in each. Long periods were found, amounting to many years, in which some months or years had over average or average rainfall but the majority had under average rainfall. However, periods of uninterrupted rainfall deficiency were regarded as more important for this study of droughts.

An index of severity was derived from amounts of rainfall deficiency in droughts of varying duration. The amount of deficiency was expressed in terms of thousandths of the

annual average rainfall for the group of stations concerned. Also, for each dry period determined from the residual mass graph, or for each period of highest intensity in a drought of long duration, a plot was made on a spot diagram of duration against amount of deficiency per month. From the spot diagram, the degree of severity (high, moderate or low) of any particular drought relative to experience in the area concerned, as shown by all plots on the diagram, could be determined and the index of severity interpreted accordingly.

In discussing effects on primary production during dry periods, frequent reference is made below to average wheat yield. Although this has tended to increase during the period under review because of improved strains, more efficient methods of cultivation, etc., the average yield is still regarded as a fairly good indicator of the severity of drought. The wheat belt is so situated that the average yield also reflects to a considerable degree the effects of drought over considerable areas of neighbouring pastoral country.

(d) *Drought Severity According to Climate.* Reference was made in the statement by Russell to the distinction between definitions of drought based on experiences in New South Wales and in the British Isles. An example of the different economic effects of droughts in a cool moist climate as compared with a dry hot climate may be seen in statistics of wheat yields. In Tasmania, since 1915, the average yield has been about 20 bushels per acre (acreage under wheat approximately 20,000 up to 1938). The lowest yields were 11.6 and 12.5 bushels per acre in wet years (1917-18 and 1916-17) and 13.4 in a dry year (1946). For Australia as a whole (acreage approximately 12 million), the average yield has been about 12 bushels per acre and the lowest 2.4 and 2.6 in years of severe drought (1902-03 and 1914-15 respectively). In terms of stock losses, the decrease in flocks and herds due to dry spells in a cool moist region with a reliable rainfall can scarcely be distinguished in statistical tables from fluctuations due to other causes. In this type of climate, droughts may involve losses of income for farmers and pastoralists as a result of poor crops, shortage of feed and water for stock and losses of stock, but it appears from production statistics that equally severe losses may occur as a result of excessively wet seasons. In the pastoral areas of inland Australia, on the other hand, cattle losses due mainly to drought may be counted by hundreds of thousands and sheep losses by millions. In Queensland, in 1947, it was reported that companies engaged in the dairying industry (in a warm moist climate with a rather high variability of rainfall) lost £2½ million due to drought in that year. In 1951, sugar losses amounted to £5 million, wheat £3 million and butter and cheese losses £1 million. Drought losses by graziers were estimated to be at least £10 million.

(e) *Major Widespread Droughts in Australia.* A brief discussion is given below of the seven major severe and widespread droughts since rainfall records commenced (those of 1864-66, 1880-86, 1888, 1895-1903, 1911-16, 1918-20 and 1939-45) and of some less severe but nevertheless widespread droughts.

1864-66 (or 68). The little data available indicate that this drought was rather severe in Victoria, South Australia, New South Wales, Queensland and Western Australia.

1880-86. Areas most affected were in Victoria (North and Gippsland), New South Wales (mainly Northern Wheat Belt, Northern Tablelands and South Coast), Queensland (1881-86 in South East with breaks, otherwise mainly in coastal areas, the Central Highlands and Central Interior in 1883-86) and South Australia (1884-86 mainly in agricultural areas).

Rainfall deficiencies in the wheat growing areas were not particularly high except in 1881 in Victoria and in the critical growth period in Western Australia and in 1885-86 in South Australia. Average wheat yields for Australia were 7.65 bushels per acre in 1880-81, 7.16 in 1881-82, 7.06 in 1882-83 and 6.03 in 1885-86. A considerable fall in sheep numbers occurred in 1884 amounting to 8 million for Australia as a whole (6 million in New South Wales and 2 million in Queensland).

1888. This occurred particularly in Victoria (North and Gippsland), Tasmania (1887-89 in South), New South Wales, Queensland (1888-89), and South Australia (central agricultural areas).

In 1888, wheat growing areas suffered a considerable rainfall deficiency in the growing season. Average yields were 3.85 bushels per acre in South Australia and 7.1 in Victoria (the principal wheat areas), 4.76 in New South Wales and 5.45 for Australia as a whole. Rainfall deficiencies were also high in much of Queensland in 1888 (including the following summer),

on the Coast of New South Wales, in Gippsland, Victoria, and in pastoral areas of South Australia. This drought however was of short duration and did not seriously affect stock numbers beyond retarding the rapid increase of that period.

1895–1903. It is difficult for most present day Australians to realise the magnitude of the effects of this drought on the economy of the country. Sheep numbers which had reached 100 million were reduced by half and cattle numbers by over 40 per cent. Average wheat yields exceeded 8 bushels per acre in only one year of the nine.

Rainfall deficiencies were very serious in South Australia, northern Victoria, New South Wales, central Australia and Tasmania where conditions were almost continuously dry from 1896 till the end of 1902. In the Western Australian agricultural areas, drought commenced in 1894 but good seasons were experienced in 1899 and 1900. In Queensland, almost continuously dry conditions prevailed from 1896, deficiencies in the wet season being very marked during 1900–1903. On the North Coast and Northern Tablelands of New South Wales, drought was intermittent until 1900 when continuous dry conditions set in until early in 1903. Areas least affected were the South Coast of New South Wales, western Victoria and the Northern Rivers district of the Northern Territory but even in these areas serious deficiencies occurred for various periods.

The lowest average wheat yield was in 1902, 2.4 bushels per acre for the Commonwealth (1.24 in New South Wales, 1.29 in Victoria and 3.64 in South Australia). In 1896, the average for Australia was 4.88 (1.66 in South Australia, 4.49 in Victoria), and in 1895, 5.19 (4.01 in Victoria, 4.2 in South Australia and 8.71 in New South Wales).

Sheep numbers had reached 106.4 million in 1892. In 1893, there was a decrease of 3.8 million, in 1895, 9.7 million, in 1897, 8 million (New South Wales 4.4 million, Queensland 1.8 million and South Australia 1.3 million), in 1898, 3.4 million (New South Wales 2.7 million), in 1899, 6.9 million (New South Wales 5 million) and in 1902, 18.4 million (New South Wales 15.2 million, Queensland 2.8 million).

Cattle numbers had reached 12.3 million by 1895. By 1898, the number had fallen to 10.8 million, by 1900 to 9.6 million and by January 1903 to 7 million.

1911–16. Rainfall deficiencies were high in the growing season of 1914 in the wheat growing areas of Victoria, New South Wales, South Australia and Western Australia, the worst drought year since 1902. The average wheat yield for the Commonwealth was only 2.58 bushels per acre (Victoria 1.38, South Australia, 1.41, New South Wales 4.65 and Western Australia 1.91).

Dry conditions in inland New South Wales extended from 1911 to 1915 or 1916 and in Western Australia from 1910 or 1911 to 1914. In Queensland and parts of the Northern Territory (Tennant Creek—Alexandria Downs), large rainfall deficiencies were experienced from 1911 to 1916, in the latter area practically continuously. In Queensland, the wet seasons failed seriously in 1912 and 1915 and on the Coast in 1916. The Northern Tablelands of New South Wales experienced similar conditions, and also the North Coast except in 1912. In Victoria and Tasmania, the drought conditions were mainly confined to the period spring 1913 to autumn 1915.

Statistics of sheep numbers showed a decline of 9.75 million in 1912 (5.9 million in New South Wales, 2 million in Victoria) and 9.35 million in 1915 (7.2 million in Queensland, 1.5 million in Victoria).

Cattle numbers decreased by nearly 2 million from 1912 to 1915 inclusive, including 1.1 million in 1915 (675,000 in Queensland, 320,000 in Victoria).

1918–20. Serious rainfall deficiencies occurred in this period in Queensland, New South Wales (except the South Coast), northern and western Victoria, South Australia, Northern Territory and Tasmania (1919–24 in the south). In 1920, there was no "wet" season in Queensland particularly on the coast. The only areas completely free from drought were the agricultural areas of Western Australia.

Average wheat yields for Australia in 1918 were 9.47 bushels per acre (New South Wales 7.6) and in 1919, 7.16 bushels per acre (New South Wales 2.98, Victoria 7.75, South Australia 7.77).

The decline in sheep numbers in 1919 was 12.4 million (9.1 million in New South Wales, 1.3 million in Victoria). Cattle numbers showed little change.

1922–23 and 1926–29. Wet season rains were deficient in Queensland in 1923 and 1926 and in coastal areas of that State in 1924. The adjacent North Coast and Tablelands of New South Wales likewise experienced dry conditions. Other areas affected by the first dry period which commenced in 1922 or earlier were inland areas of New South Wales, pastoral areas of South Australia, northern and western Victoria and part of the north-west of Western Australia.

The average wheat yield in New South Wales was 9.74 bushels per acre in 1922–23 and that for the Commonwealth 11.21 bushels.

On the Central Coast of Queensland, very dry conditions persisted until 1926. In inland areas of New South Wales and in Victoria, another dry period was experienced from 1925–26 until early 1930. This included the cereal growing seasons of 1927 and 1928. In the Western and Gippsland Districts of Victoria, however, the dry conditions ended in 1927. Somewhat similar conditions prevailed in Tasmania.

In the inland areas of Queensland, an even more severe drought period was experienced from 1925 to 1931 in the southern interior and until 1935 in the central interior. The year 1926 was very bad in Queensland. In the Northern Territory, a dry period commenced in 1925 or earlier and continued until 1929 with large rainfall deficiencies. Similar conditions prevailed in pastoral areas of South Australia from the end of 1926 until 1930. In parts of the north west of Western Australia, there were heavy rainfall deficiencies from 1922 to 1927.

The average wheat yield for the Commonwealth in 1927–28 was 9.63 bushels per acre (8.92 in New South Wales, 8.54 in Victoria and 8.16 in South Australia).

A fall of nearly 4 million in sheep numbers occurred in Queensland in 1926 but this was more than offset by increases in the other States. In 1927, however, there was a fall of 3.4 million in Commonwealth numbers (5.2 million in New South Wales partly offset by rises in other States). Cattle numbers fell by 1.4 million in 1926 (1 million in Queensland).

1933–38. During the thirties, there were frequent dry periods of some severity in some parts of the Commonwealth, the most notable up to 1935 being in Queensland in 1931 and 1932–33. In 1935, the wet season again failed in Queensland and to a lesser degree also in parts of the Northern Territory and the north-west of Western Australia. In the latter region, a long dry period commenced in 1934–1935 and persisted until 1941. In Western Australian agricultural areas, the period 1934–38 was persistently dry. Victoria and inland areas of New South Wales experienced dry conditions in 1937–38 and Tasmania at intervals in 1935–37 (1933 and the early part of 1934 having also been dry).

Tropical areas experiencing dry conditions in 1937–38 included Queensland and the Northern Territory. Rainfall deficiencies were considerable in southern inland areas of Queensland and also in western New South Wales. Rather dry conditions persisted throughout most of the period 1935–39 on the coast of New South Wales while on the Northern Tablelands there was a long period of under average rainfall from 1935 to 1940 inclusive.

Except in 1937 and 1938, most of the cereal growing areas were free from drought and yields generally were fairly satisfactory. In 1934, the average wheat yield for the Commonwealth was 10.63 bushels per acre (8.61 in South Australia) and in 1938, 10.83 bushels per acre (Victoria 6.59). Sheep numbers decreased by 3 million in 1933 (1.5 million in New South Wales, 1.3 million in Queensland), 4.2 million in 1935 (1.4 million in New

South Wales, 3.5 million in Queensland), and 2.3 million in 1938 (2.7 million in New South Wales offset by increases elsewhere). In Western Australia, there was a decrease of 2.3 million in 1936 and 1937 but Commonwealth numbers increased in these years. Cattle numbers decreased by about 1.2 million between 1935 and 1939 (670,000 in New South Wales and about 400,000 in Victoria).

1939-45. The main dry periods in these years were 1940 and 1943-45. In 1940, serious deficiencies occurred in the cereal growing season in Victoria, inland New South Wales, South Australia and Western Australia. The average wheat yield for the Commonwealth was reduced to 6.5 bushels per acre (5.06 in Victoria, 5.37 in New South Wales, 6.97 in South Australia and 8.02 in Western Australia). In 1944, serious deficiencies again occurred in these areas resulting in an average wheat yield of 6.25 bushels per acre for the Commonwealth (1.63 in Victoria, 5.7 in South Australia and 6.02 in New South Wales).

Persistent rainfall deficiencies occurred throughout most of the period 1939-1945 in pastoral areas of South Australia, western New South Wales, southern Queensland (1940-47) and on the coast of New South Wales. The serious drought conditions in the De Grey, Fortescue, Gascoyne and Murchison districts of Western Australia in the mid thirties continued until 1941. In Tasmania, the period as a whole was one of rainfall deficiency, 1945 being an exceptionally dry year in the south.

Sheep numbers in Australia had reached 125 million by 1942. In the next four years, there was a decline of 29 million. Approximately 18 million occurred in 1944 (New South Wales 10 million, Victoria 3 million, Queensland 2 million, South Australia 2 million and Western Australia 1 million), and 9 million in 1945 (2.6 million in New South Wales, 2.4 million in Queensland and 1.8 million in Victoria).

1946-49. The summers of 1947, 1948 and 1949 were marked by considerable rainfall deficiencies in Queensland and the Northern Territory and the cereal growing season of 1946 in inland areas of New South Wales. The wet seasons of 1948 and 1949 were also deficient in rainfall in parts of the north-west of Western Australia as were the cereal growing periods in the agricultural areas of that State. The wheat yield for 1946 was 8.9 bushels per acre for the Commonwealth (3.5 in New South Wales, 2.8 in Queensland and 9.81 in Western Australia), but in other years was very satisfactory.

1951-55. The wet season of 1951-52 was marked by serious rainfall deficiencies in Queensland, the Northern Territory and the north-west of Western Australia. In Queensland and the Northern Territory, the dry conditions set in in the late summer of 1951. In the Kimberley division of Western Australia, serious deficiencies persisted until about March 1954 and other parts of that State experienced prolonged periods of rainfall deficiency in these years. Growing seasons for cereals were on the whole free from serious drought and wheat yields were maintained at a high level. In the 1952-53 season, the average for the Commonwealth, 19.1 bushels per acre, was the highest on record, the total yield being 195 million bushels. The average yield for 1953-54, 18.4 bushels per acre, was also higher than any prior to 1952-53. The highest wheat yield on record was 220 million bushels in 1947-48 from 13.9 million acres. Acreages in 1952-53 and 1953-54 were respectively 10.6 and 11.2 million, there having been a decline since the early thirties (18 million in 1930-31).

Drought in tropical Australia in 1951-52 resulted in heavy cattle and sheep losses, though this is not very apparent from statistics for the States and Commonwealth as a whole. Sheep numbers in Queensland and New South Wales showed a decline of $1\frac{1}{2}$ million in the year ended March 1952, but for the Commonwealth there was an increase of about $1\frac{1}{2}$ million. Cattle numbers decreased by about 300,000.

It is of interest to note that the present succession of good agricultural seasons, at least in the wheat belt, has been the longest on record with average wheat yields of over 15 bushels per acre. This is presumably due in some measure to improvements in wheat strains and more efficient farming methods. A similar period with average yields above 10 bushels per acre for the Commonwealth was the decade 1930-39. In these years, in New South Wales, average yields were 12 to 17 bushels per acre. Victoria experienced a poor year in 1938 and South Australia had only moderate yields in 1930, 1933, 1934 and 1936 while in Western Australia the yields in the three years 1934-36 were below 10 bushels per acre.

(f) *Conclusions.* The conclusion to be drawn from this study is that, during the last 100 years, Australia has been subjected to at least seven major droughts affecting the greater part of the continent and several other droughts causing severe losses in restricted areas. The drought of 1895 to 1903 was undoubtedly the most disastrous in its effects on primary producers.

The method of tracing fluctuations of rainfall, month by month, in relation to the normal demonstrates that in all parts of the Commonwealth large and irregular variations in rainfall occur and there is no reason to doubt that great fluctuations, especially in inland areas, will be repeated in the future. It may also be argued that the worst conditions experienced in the last 100 years do not represent the worst conditions ever likely to be experienced. Although details of experiences of primary producers and other settlers during drought periods make a gloomy picture, they serve to indicate the nature of measures previously adopted to counteract the worst effects of drought and which are capable of further expansion in the future. These include conservation of fodder in good seasons and increased facilities for the storage of fodder and grain, the provision of adequate transport facilities for the economic removal of starving stock from, and the supply of fodder and water to, drought stricken areas and the expansion of irrigation to areas where this is practicable. The rapid expansion of primary industries in Australia in the last fifty years has greatly increased the vulnerability of the economic position of the country to drought. However, the fact that settlement has been maintained and has prospered in many areas subject to frequent and severe droughts indicates that the rich returns of good seasons, if carefully husbanded, are sufficient to tide over lean periods.

Residual mass graphs show that the incidence of droughts shows no regular rhythm in time of onset, duration or extent of territory affected, indicating that there is little or no prospect of successfully forecasting drought from an assumed occurrence of rainfall cycles. The little work done on the study of synoptic weather types in drought periods suggests that droughts are related to certain characteristics of the general circulation of the atmosphere including the absence of favourable atmospheric flow patterns and of an effective rain producing mechanism in such periods over areas affected by drought.

Results of upper air observations which have become available in recent years show that in wet years the frequent large scale development of upper cyclones ("cut off lows") and troughs is a striking feature while in dry periods they are almost entirely absent. The shift of this form of activity in the general circulation of the atmosphere from one region of the earth's surface to another is a subject for world wide study.

10. *Barometric Pressures.*—The mean annual barometric pressure (corrected to sea level and standard gravity) in Australia varies from 29.80 inches on the north coast to 29.92 inches over the central and 30.03 inches in the southern parts of the continent. In January, the mean pressure ranges from 29.70 inches in the northern and central areas to 29.95 inches in the southern. The July mean pressure ranges from 29.90 inches at Darwin to 30.12 inches at Alice Springs. Barometer readings corrected to mean sea level and standard gravity have, under anticyclonic conditions, ranged as high as 30.935 inches (at Hobart on 13th July, 1846) and have fallen as low as 27.55 inches. This record low was registered at Mackay during a tropical hurricane on 21st January, 1918. An almost equally abnormal reading of 27.88 inches was recorded at Innisfail during a similar storm on 10th March, 1918. For graphs of Mean Barometric Pressure at Capital Cities, see Official Year Book No. 37, page 35.

11. *Wind.*—(i) *Trade Winds.* The two distinctive wind currents in Australia are, as previously stated, the south-east trade and the "prevailing" westerly winds. As the belt of the earth's atmosphere in which they blow apparently follows the sun's ecliptic path north and south of the equator, so the area of the continent affected by these winds varies at different seasons of the year. During the summer months, the anticyclonic belt travels in high latitudes, thereby bringing the south-east trade winds as far south

as 30° south latitude. The "prevailing" westerly winds retreat a considerable distance to the south of Australia, and are less in evidence in the hot months. When the sun passes to the north of the equator, the south-east trade winds follow it, and only operate to the north of the tropics for the greater part of the winter. The westerly winds come into lower latitudes during the same period of the year. They sweep across the southern areas of the continent from Cape Leeuwin to Cape Howe, and during some seasons are remarkably persistent and strong, and occasionally penetrate to almost tropical latitudes.

(ii) *North-west Monsoon.* As the belt of south-east trade winds retreats southward during the summer, it is replaced in the north and north-west of Australia first by a sequence of light variable winds and then by the north-west monsoon. In Australia, the north-west monsoon has not the persistence nor regularity of the Indian south-west monsoon but is sufficiently characteristic for the summer in the north of Australia to be called the "North-west Season". In central and eastern Queensland, the north-west monsoon in the summer has comparatively little effect and the trade winds, albeit weakened, are still dominant winds. With the migration of the sun northward in the autumn, the north-west monsoon is replaced first by light variable winds and then by the trade winds.

Further particulars of Australian wind conditions and meteorology will be found in Official Year Book No. 38, pages 58-61. Reference should also be made to the wind rose diagrams on pages 43 and 44.

(iii) *Cyclones and Storms.* The "elements" in Australia are ordinarily peaceful, and while destructive cyclones have visited various parts, more especially coastal areas, such visitations are rare, and may be properly described as erratic.

During the winter months, the southern shores of the continent are subject to deep depressions of the southern low-pressure belt. They are felt most severely over the south-western parts of Western Australia, to the south-east of South Australia, in Bass Strait, along the coastline of Victoria, and on the west coast of Tasmania. Apparently the more violent wind pressures from these disturbances are experienced in their northern half, or in that part of them which has a north-westerly to a south-westerly circulation.

The north-east coast of Queensland is occasionally visited by hurricanes from the north-east tropics. During the first four months of the year, these hurricanes appear to have their origin in the neighbourhood of the South Pacific Islands, their path being a parabolic curve first to the south-west and finally towards the south-east.

Very severe cyclones, locally known as "willy willies," are peculiar to the north-west coast of Western Australia from the months of November to April, inclusive. They usually originate over the ocean to the north or north-west of Australia, and travel in a south-westerly direction with continually increasing force, displaying their greatest energy near Cossack and Onslow, between latitudes 20° and 22° South. The winds in these storms, like those from the north-east tropics, are very violent and destructive. The greatest velocities are usually to be found in the south-eastern quadrant of the cyclones, with north-east to east winds. After leaving the north-west coast, these storms either travel southwards, following the coast-line, or cross the continent to the Great Australian Bight. When they take the latter course, their track is marked by torrential rains, as much as 29.41 inches, for example, being recorded in 24 hours at Whim Creek from one such occurrence. Falls of 10 inches and over have frequently been recorded in the northern interior of Western Australia from similar storms.

Some further notes on severe cyclones and on "southerly bursters", a characteristic feature of the eastern part of Australia, appeared in early issues of the Official Year Book (see No. 6, pp. 84-86), and a special article dealing with "Australian Hurricanes and Related Storms" appeared in Official Year Book No. 16, pp. 80-84.

Depressions vary considerably in their isobaric forms, intensity and other characteristics. Some bring rain in variable quantities, some heat and others mainly wind. A common type in southern Australia is the "Λ" shaped trough with an abrupt "backing" of the wind or "line squall" as it passes. The cold front is most frequently found through the centre of the "trough" because it is along this line, and extending into the upper levels of the atmosphere, that the demarcation of different air masses is so well defined. The best rains in inland Australia occur when extensive masses of warm moist tropical air move into the interior and are forced to rise by convergence of flow or by impact with a cold air stream.

The speed of low pressure systems is very variable, but in general in southern latitudes the movement is of the order of 500 to 700 miles per day.

12. **Influences affecting Australian Climate.**—(i) *General.* Australian history does not cover a sufficient period, nor is the country sufficiently occupied, to ascertain whether or not the advance of settlement has materially affected the climate as a whole. Local changes have, however, taken place, a fact which suggests that settlement and the treatment of the land have a distinct effect on local conditions. For example, low-lying lands on the north coast of New South Wales, which originally were seldom subject to frosts, have, with the deforestation of the surrounding hills, experienced annual visitations, the probable explanation being that through the absence of trees the cold air of the highlands now flows unchecked and untempered down the sides of the hills to the valleys and lower lands.

(ii) *Influence of Forests on Climate.* As already indicated, forests doubtless exercise a great influence on local climate, and hence, to the extent that forestal undertakings will allow, the weather can be controlled by human agency. The direct action of forests is an equalizing one; thus, especially in equatorial regions, and during the warmest portion of the year, they considerably reduce the mean temperature of the air. They also reduce the diurnal extremes of shade temperatures by altering the extent of radiating surface by evaporation, and by checking the movement of air, and while decreasing evaporation from the ground, they increase the relative humidity. Vegetation greatly diminishes the rate of flow-off of rain and the washing away of surface soil, and when a region is protected by trees a steadier water supply is ensured, and the rainfall is better conserved. In regions of snowfall, the supply of water to rivers is similarly regulated, and without this and the sheltering influence of ravines and "gullies," watercourses supplied mainly by melting snow would be subject to alternate periods of flooding and dryness. This is borne out in the case of the inland rivers, the River Murray, for example, which has never been known to become dry, deriving its steadiness of flow mainly through the causes indicated.

(iii) *Direct Influence of Forests on Rainfall.* Whether forests have a direct influence on rainfall is a debatable question, some authorities alleging that precipitation is undoubtedly induced by forests, while others take the opposite view.

Sufficient evidence exists, however, to prove that, even if the rainfall is not increased, the beneficial climatic effect of forest lands more than warrants their protection and extension. Rapid rate of evaporation, induced by both hot and cold winds, injures crops and makes life uncomfortable on the plains, and, while it may be doubted that the forest aids in increasing precipitation, it must be admitted that it does check winds and the rapid evaporation due to them. Trees as wind-breaks have been successfully planted in central parts of the United States of America, and there is no reason why similar experiments should not be successful in many parts of the treeless interior of Australia. The belts should be planted at right angles to the direction of the prevailing parching winds, and if not more than half a mile apart will afford shelter to the enclosed areas.

13. **Rainfall and Temperatures, Various Cities.**—Official Year Book No. 34, page 28, shows rainfall and temperature and No. 38, page 42, temperature, for various important cities throughout the world and for the Australian capitals.

14. **Climatological Tables.**—The averages and extremes for a number of climatological elements, which have been determined from long series of observations at the Australian capitals up to and including the year 1957, are given on the following eight pages.

NOTE.—The following points apply throughout:—

- (i) Where records are available, mean or average values have been calculated on a standard period of 30 years from 1911 to 1940.
- (ii) Extreme values have been extracted from all available years of actual record, but the number of years quoted does not include intervening periods when observations were temporarily discontinued.

CLIMATOLOGICAL DATA: PERTH, WESTERN AUSTRALIA.
(LAT. 31° 57' S., LONG. 115° 51' E. HEIGHT ABOVE M.S.L. 210 FT.)
Barometer, Wind, Evaporation, Lightning, Clouds and Clear Days.

Month.	Bar. corrected to 32° F. M.n. Sea Level and Standard Gravity from 9 a.m. and 3 p.m. readings.	Wind. (Height of Anemometer 71 feet.)					Mean Amount of Evaporation (inches).	No. of Days of Lightning.	Mean Amount of Clouds, 9 a.m., 3 p.m., 9 p.m.(d)	No. of Clear Days.
		Average Miles per Hour.	Highest Mean Speed in One Day (miles per hour).	Highest Gust Speed (miles per hour).	Prevailing Direction.					
					9 a.m.	3 p.m.				
No. of years of observations.	30(b)	30(b)	59	45	30(b)	30(b)	30(b)	30(b)	30(b)	
January ..	29.897	10.1	33.2 27/98	49	E	SSW	10.37	2	2.9	14
February ..	29.922	9.9	27.1 6/08	54	ENE	SSW	8.63	2	3.1	13
March ..	29.976	9.4	27.1 6/13	66	E	SSW	7.52	2	3.5	12
April ..	30.071	7.9	39.8 25/00	61	ENE	SSW	4.62	2	4.2	9
May ..	30.062	7.8	34.4 29/32	73	NE	WSW	2.80	3	5.4	6
June ..	30.068	7.8	38.1 17/27	80	N	NW	1.82	2	5.9	5
July ..	30.082	8.2	42.3 20/26	73	NNE	W	1.76	2	5.6	5
August ..	30.084	8.7	40.3 15/03	77	N	WNW	2.37	2	5.6	6
September ..	30.073	8.7	36.0 11/05	75	ENE	SSW	3.44	1	4.9	8
October ..	30.033	9.3	33.7 6/16	65	SE	SW	5.38	1	4.8	8
November ..	29.989	9.9	32.4 18/97	63	E	SW	7.65	2	3.9	9
December ..	29.923	10.2	32.3 6/22	64	E	SSW	9.69	2	3.2	13
Year { Totals ..	—	—	—	—	—	—	66.05	23	—	108
Year { Averages ..	30.015	9.0	—	—	E	SSW	—	—	4.4	—
Year { Extremes ..	—	—	42.3 20/7/26	80	—	—	—	—	—	—

(a) Scale 0-10. (b) Standard 30 years' normal (1911-1940).

Temperature and Sunshine.

Month.	Mean Temperature (°Fahr.).			Extreme Shade Temperature (°Fahr.).			Extreme Temperature (°Fahr.).			Mean Daily Hours of Sunshine.
	Mean Max.	Mean Min.	Mean	Highest.	Lowest.	Extreme Range.	Highest in Sun.	Lowest on Grass.		
									No. of years over which observation extends.	
No. of years over which observation extends.	30(a)	30(a)	30(a)	61	61	61	53	59	30(a)	
January ..	84.6	63.3	73.9	110.7 29/56	48.6 20/25	62.1	177.3 22/14	39.5 20/25	10.4	
February ..	85.1	63.5	74.3	112.2 8/33	47.7 1/02	64.5	173.7 4/34	39.8 1/13	9.8	
March ..	81.3	61.5	71.4	106.4 14/22	45.8 8/03	60.6	175.9 23/47	36.7 8/03	8.8	
April ..	76.3	57.4	66.8	99.7 9/10	39.3 20/14	60.4	157.0 8/16	31.0 20/14	7.5	
May ..	69.0	52.8	60.9	90.4 2/07	34.3 11/14	56.1	146.0 4/25	25.3 11/14	5.7	
June ..	64.4	49.8	57.1	81.7 2/14	34.9 22/55	46.8	135.5 9/14	26.3 11/37	4.8	
July ..	62.8	48.0	55.4	76.4 21/21	34.2 7/16	42.2	133.2 13/15	25.1 30/20	5.4	
August ..	63.8	48.4	56.1	82.0 21/40	35.3 31/08	46.7	145.1 29/21	26.7 24/35	6.0	
September ..	66.8	50.4	58.6	90.9 30/18	36.7 22/56	54.2	153.6 29/16	27.2 (b)	7.2	
October ..	69.7	52.6	61.1	95.3 30/22	40.0 16/31	55.3	157.5 31/36	29.8 16/31	8.1	
November ..	76.7	57.3	67.0	104.6 24/13	42.0 1/04	62.6	167.0 30/25	35.5 (c)	9.6	
December ..	81.2	60.9	71.0	107.9 20/04	47.5 29/57	60.4	168.8 11/27	38.0 29/5 7	10.4	
Year { Averages ..	73.5	55.5	64.5	—	—	—	—	—	7.8	
Year { Extremes ..	—	—	—	112.2 8/2/33	34.2 7/7/16	78.0	177.3 22/1/14	25.1 30/7/20	—	

(a) Standard 30 years' normal (1911-1940).

(b) 8/52 and 6/56.

(c) 6/10 and 14/12.

Humidity, Rainfall and Fog.

Month.	Vapour Pressure (inches)	Rel. Hum. (%) at 9 a.m.			Rainfall (inches).				Fog.	
		Mean.	Highest Mean.	Lowest Mean.	Mean Monthly.	Mean No. of Days of Rain.	Greatest Monthly.	Least Monthly.		
										No. of years over which observation extends.
No. of years over which observation extends.	30(a)	30(a)	59	59	30(a)	30(a)	82	82	82	30(a)
January ..	0.438	51	61	41	0.33	3	2.17 1879	Nil (b)	1.74 27/79	0
February ..	0.434	51	65	43	0.50	3	6.55 1955	Nil (b)	3.43 17/55	0
March ..	0.432	57	66	46	0.90	5	5.71 1934	Nil (b)	3.03 9/34	0
April ..	0.397	61	73	51	1.75	8	5.85 1926	Nil 1920	2.62 30/04	1
May ..	0.365	70	81	61	5.14	15	12.13 1879	0.98 1903	3.00 17/42	2
June ..	0.337	75	83	68	7.55	17	18.75 1945	2.16 1877	3.90 10/20	2
July ..	0.322	76	84	69	7.08	19	12.28 1926	2.42 1876	3.00 4/91	2
August ..	0.316	71	83	62	5.78	19	12.53 1945	0.46 1902	2.91 14/45	1
September ..	0.341	66	75	58	3.37	15	7.84 1923	0.34 1916	1.82 4/31	0
October ..	0.345	60	75	52	2.30	12	7.87 1890	0.15 1946	1.73 3/33	0
November ..	0.374	52	63	41	0.75	7	2.78 1916	Nil 1891	1.40 15/48	0
December ..	0.409	51	63	44	0.54	5	3.05 1888	Nil (c)	1.72 1/88	0
Year { Totals ..	—	—	—	—	35.99	128	—	—	—	8
Year { Averages ..	0.370	62	—	—	—	—	—	—	—	—
Year { Extremes ..	—	—	84	41	—	—	18.75 6/1945	Nil (d)	3.90 10/6/20	—

(a) Standard 30 years' normal (1911-1940).

(b) Various years.

(c) 1886 and 1924.

(d) November to April, various years.

CLIMATOLOGICAL DATA; DARWIN, NORTHERN TERRITORY.
(LAT. 12° 28' S., LONG. 130° 51' E. HEIGHT ABOVE M.S.L. 97 Ft.)
Barometer, Wind, Evaporation, Lightning, Clouds and Clear Days.

Month.	Bar. corrected to 32° F. M.S. Sea Level and Standard Gravity from 9 a.m. and 3 p.m. readings.	Wind.					Mean Amount of Evaporation (inches).	No. of Days of Lightning.	Mean Amount of Clouds, 9 a.m., 3 p.m., 9 p.m. (a)	No. of Clear Days.
		Average Miles per Hour.	Highest Mean Speed in One Day (miles per hour).	Highest Gust Speed (miles per hour).	Prevailing Direction.					
					9 a.m.	3 p.m.				
No. of years of observations.	30	14	—	—	—	—	—	30	30	30
January	29.706	6.1	—	—	—	NW & S	W & NW	16	7.1	1
February	29.728	6.7	—	—	—	W & S	W & NW	16	7.0	1
March	29.751	5.3	—	—	—	SE	W & NW	14	6.2	3
April	29.809	6.1	—	—	—	SE	E	6	3.5	11
May	29.859	6.5	—	—	—	SE	E	1	2.1	19
June	29.892	6.5	—	—	—	SE	E & SE	0	1.6	22
July	29.911	6.2	—	—	—	SE	E & SE	0	1.4	23
August	29.914	5.9	—	—	—	SE	NW & N	0	1.3	23
September	29.886	6.2	—	—	—	SE & S	NW & N	1	2.0	18
October	29.850	6.2	—	—	—	S	NW & N	8	3.2	10
November	29.797	5.5	—	—	—	W & S	NW & N	17	4.8	4
December	29.738	6.2	—	—	—	NW & S	NW & N	17	6.0	2
Year { Totals ..	29.820	6.1	—	—	—	SE	NW	96	3.9	137
Year { Averages ..	—	—	—	—	—	—	—	—	—	—
Year { Extremes ..	—	—	—	—	—	—	—	—	—	—

(a) Scale 0-10.

Temperature and Sunshine.

Month.	Mean Temperature (°Fahr.).			Extreme Shade Temperature (°Fahr.).		Extreme Range.	Extreme Temperature (°Fahr.).		Mean Daily Hours of Sunshine.	
	Mean Max.	Mean Min.	Mean	Highest.	Lowest.		Highest in Sun.	Lowest on Grass.		
										No. of years over which observation extends.
January	89.9	77.3	83.6	99.1	8/28	69.2	21/44	168.0	26/42	—
February	89.8	77.1	83.4	97.0	13/37	63.0	25/49	163.6	23/38	—
March	90.2	77.1	83.6	100.0	8/31	66.6	31/45	165.6	23/38	—
April	91.9	75.9	83.9	98.0	19/24	60.8	11/43	163.0	1/38	—
May	90.9	72.6	81.4	96.8	(b)	59.2	8/49	160.0	5/20	—
June	87.5	69.5	78.5	98.6	17/37	55.3	18/49	155.2	2/16	—
July	86.6	67.8	77.2	94.0	16/21	50.7	29/42	156.0	28/17	—
August	88.5	69.7	79.1	96.0	30/36	57.0	16/57	152.2	28/16	—
September	91.0	73.9	82.5	99.0	25/28	63.8	1/46	157.0	(c)	—
October	92.6	77.2	84.9	99.0	14/33	68.5	26/45	160.5	30/38	—
November	93.2	78.2	85.7	101.0	27/24	67.4	12/45	170.4	14/37	—
December	92.0	78.1	85.0	100.4	13/31	68.5	24/41	169.0	26/23	—
Year { Averages ..	90.3	74.5	82.4	—	—	—	—	170.4	—	—
Year { Extremes ..	—	—	—	101.0	27/11/24	50.7	29/7/42	—	14/11/37	—

(a) Years 1918-41 at Post Office, 1942-56 at aerodrome; sites not strictly comparable.
(c) 28/16 and 3/21.

(b) 2/37 and 2/42.

Humidity, Rainfall and Fog.

Month.	Vapour Pressure (inches)	Rel. Hum. (%) at 9 a.m.			Rainfall (inches).					Fog. Mean No. of Days of Fog.			
		Mean.	Highest Mean.	Lowest Mean.	Mean Monthly.	Mean No. of Days of Rain.	Greatest Monthly.	Least Monthly.	Greatest in One Day.				
											No. of years over which observation extends.	57	57
January	0.925	78	89	69	16.18	20	27.86	1906	2.25	1930	11.67	7/97	0.0
February	0.920	79	88	71	12.37	18	25.74	1955	0.44	1931	5.25	15/49	0.0
March	0.912	78	84	69	11.18	17	21.88	1898	0.81	1911	7.18	6/19	0.0
April	0.800	69	80	60	3.08	6	23.74	1891	Nil	(a)	5.51	1/29	0.0
May	0.652	63	76	49	0.33	1	14.00	1953	Nil	(a)	2.19	6/22	0.0
June	0.545	61	75	52	0.09	1	1.53	1902	Nil	(a)	1.32	10/02	0.4
July	0.522	59	71	47	0.01	0	2.56	1900	Nil	(a)	1.71	2/00	1.1
August	0.613	63	73	53	0.02	0	3.00	1870	Nil	(a)	1.06	14/09	0.7
September	0.732	65	73	54	0.60	2	2.72	1950	Nil	(a)	2.00	26/50	0.2
October	0.832	65	72	60	1.93	5	13.34	1954	Nil	(a)	3.74	18/56	0.0
November	0.868	68	75	62	4.32	10	15.72	1938	0.40	1870	4.73	9/51	0.0
December	0.890	73	83	65	8.57	15	22.38	1910	0.98	1934	7.87	28/10	0.0
Year { Totals ..	—	—	—	—	58.68	95	—	—	—	—	—	—	2.4
Year { Averages ..	0.764	68	—	—	—	—	—	—	—	—	—	—	—
Year { Extremes ..	—	—	89	47	—	—	27.86	1/06	Nil	(b)	11.67	7/1197	—

(a) Various years.

(b) April to October, various years.

CLIMATOLOGICAL DATA; ADELAIDE, SOUTH AUSTRALIA.
 (LAT. 34° 56' S., LONG. 138° 35' E. HEIGHT ABOVE M.S.L. 140 FT.)
 Barometer, Wind, Evaporation, Lightning, Clouds and Clear Days.

Month.	Bar. corrected to 32° F., Mm. Sea Level and Standard Gravity from 9 a.m. and 3 p.m. readings.	Wind. (Height of Anemometer 75 feet.)					Mean Amount of Evaporation (inches).	No. of Days of Lightning.	Mean Amount of Clouds, 9 a.m. to 3 p.m., 9 p.m. to a.m.	No. of Clear Days.	
		Average Miles per Hour.	Highest Mean Speed in One Day (miles per hour).	Highest Gust Speed (miles per hour).	Prevailing Direction.						
					9 a.m.	3 p.m.					
No. of years of observations.	30(b)	30(b)	80	41	30(b)	30(b)	30(b)	30(b)	30(b)		
January ..	29.917	9.9	31.6	19/99	72	SW	SW	9.27	2.3	3.6	12.9
February ..	29.953	8.8	28.8	22/96	64	NE	SW	7.56	2.0	3.7	11.2
March ..	30.037	8.3	26.2	9/12	67	S	SW	6.39	1.8	4.0	10.6
April ..	30.119	8.0	32.2	10/96	81	NE	SW	3.78	1.5	5.2	7.2
May ..	30.131	8.1	31.7	9/80	67	NE	NW	2.27	1.3	5.8	4.9
June ..	30.119	8.3	31.3	12/78	67	NE	N	1.37	1.3	6.1	4.1
July ..	30.111	8.5	28.1	25/82	60	NE	NW	1.34	1.5	6.0	4.3
August ..	30.084	9.2	32.2	31/97	57	NE	SW	1.99	2.0	5.5	5.6
September ..	30.050	9.2	30.0	2/87	69	NNE	SW	3.05	2.0	5.3	5.8
October ..	30.007	9.8	32.0	28/98	73	NNE	SW	5.03	2.8	5.3	5.7
November ..	29.990	9.9	32.2	7/48	79	SW	SW	6.89	3.3	4.9	7.2
December ..	29.922	9.9	28.1	12/91	75	SW	SW	8.74	2.2	4.2	9.5
Year { Totals ..	—	—	—	—	—	—	—	57.68	24.0	—	89.0
Year { Averages ..	30.037	9.0	—	—	—	NE	SW	—	—	5.0	—
Year { Extremes ..	—	—	32.2 (c)	81	—	—	—	—	—	—	—

(a) Scale 0-10. (b) Standard 30 years' normal (1911-1940). (c) 10/4/1896, 31/8/1897 and 7/11/1948.

Temperature and Sunshine.

Month.	Mean Temperature (°Fahr.).			Extreme Shade Temperature (°Fahr.).		Extreme Range.	Extreme Temperature (°Fahr.).		Mean Daily Hours of Sunshine.					
	Mean Max.	Mean Min.	Mean	Highest.	Lowest.		Highest in Sun.	Lowest on Grass.						
										No. of years over which observation extends.	101	101	101	54(b)
January ..	84.8	61.0	72.9	117.7	12/39	45.1	21/84	72.6	180.0	18/82	36.5	14/79	10.0	
February ..	85.7	61.8	73.7	113.6	12/99	45.5	23/18	68.1	170.5	10/00	35.8	23/26	9.3	
March ..	81.3	59.1	70.2	110.5	9/34	43.9	21/33	66.6	174.0	17/83	32.1	21/33	7.9	
April ..	73.0	54.4	63.7	98.6	5/38	39.6	15/59	59.0	155.0	1/83	30.2	16/17	6.0	
May ..	66.8	50.8	58.8	89.5	4/21	36.9	26/95	52.6	148.2	12/79	25.6	19/28	4.8	
June ..	61.0	46.6	53.8	76.0	23/65	32.5	(c)	43.5	138.8	18/79	21.0	24/44	4.2	
July ..	59.9	45.4	52.7	74.0	11/06	32.0	24/08	42.0	134.5	26/90	22.1	30/29	4.3	
August ..	62.3	46.2	54.3	85.0	31/11	32.3	17/59	52.7	140.0	31/92	22.8	11/29	5.4	
September ..	66.8	48.3	57.5	91.3	29/44	32.7	4/58	58.6	160.5	23/82	25.0	25/27	6.3	
October ..	72.5	51.7	62.1	102.9	21/22	36.0	—/57	66.9	162.0	30/21	27.8	(d)	7.3	
November ..	78.1	55.4	66.7	113.5	21/65	40.8	2/09	72.7	166.9	20/78	31.5	2/09	8.6	
December ..	82.6	58.9	70.7	114.6	29/31	43.0	(e)	71.6	175.7	7/99	32.0	4/84	9.5	
Year { Averages ..	72.9	53.3	63.1	—	—	—	—	—	—	—	—	—	—	7.0
Year { Extremes ..	—	—	—	117.7	12/11/39	32.0	24/7/08	85.7	180.0	18/11/82	21.0	24/6/44	—	—

(a) Standard 30 years' normal (1911-1940). (b) Records incomplete, 1931-34. Discontinued, 1934.
 (c) 27/1876 and 24/1944. (d) 4/1931 and 2/1918. (e) 16/1861 and 4/1906.

Humidity, Rainfall and Fog.

Month.	Vapour Pressure (inches).	Rel. Hum. (%) at 9 a.m.			Rainfall (inches).					Fog. Mean No. of Days of Fog.			
		Mean 9 a.m.	Highest Mean.	Lowest Mean.	Mean Monthly.	Mean No. of Days of Rain.	Greatest Monthly.	Least Monthly.	Greatest in One Day.				
											No. of years over which observation extends.	30(a)	89
January ..	0.327	39	59	29	0.76	5	4.00	1850	Nil (b)	2.30	2/89	0.0	
February ..	0.352	41	56	30	1.10	5	6.09	1925	Nil (b)	5.57	7/25	0.0	
March ..	0.332	44	58	29	0.87	5	4.60	1878	Nil (b)	3.50	5/78	0.0	
April ..	0.329	55	72	37	1.45	10	6.78	1853	Nil 1945	3.15	5/60	0.0	
May ..	0.313	64	76	49	2.49	13	7.75	1875	0.10	1934	2.75	1/53	0.6
June ..	0.294	75	84	67	2.93	15	8.58	1916	0.42	1886	2.11	1/20	1.1
July ..	0.282	75	87	66	2.49	16	5.38	1865	0.37	1899	1.75	10/65	1.4
August ..	0.282	68	78	54	2.58	16	6.24	1852	0.33	1944	2.23	19/51	0.4
September ..	0.289	59	72	44	2.39	13	5.83	1923	0.27	1951	1.59	20/23	0.2
October ..	0.287	48	67	29	1.54	10	4.38	1948	0.17	1914	2.24	16/08	0.0
November ..	0.292	41	57	31	1.22	8	4.10	1934	0.04	1885	2.08	7/34	0.0
December ..	0.322	40	50	31	1.27	6	3.98	1861	Nil	1904	2.42	23/13	0.0
Year { Totals ..	—	—	—	—	21.09	122	—	—	—	—	—	—	3.7
Year { Averages ..	0.304	52	—	—	—	—	—	—	—	—	—	—	—
Year { Extremes ..	—	—	87	29	—	—	8.58	6/1916	Nil (c)	5.57	7/2/25	—	—

(a) Standard 30 years' normal (1911-1940). (b) Various years. (c) December to April, various years.

CLIMATOLOGICAL DATA; BRISBANE, QUEENSLAND.
(LAT. 27° 28' S., LONG. 153° 2' E. HEIGHT ABOVE M.S.L. 134 FT.)
Barometer, Wind, Evaporation, Lightning, Clouds and Clear Days.

Month.	Bar. corrected to 32° F. M. Sea Level and Standard Gravity from 9 a.m. and 3 p.m. readings.	Wind. (Height of Anemometer 105 feet.)						Mean Amount of Evaporation (inches).	No. of Days of Lightning.	Mean Amount of Clouds, 9 a.m., 3 p.m., 9 p.m.(a)	No. of Clear Days.
		Average Miles per Hour.	Highest Mean Speed in One Day (miles per hour).	Highest Gust Speed (miles per hour).	Prevailing Direction.						
					9 a.m.	3 p.m.					
No. of years of observations.	30(b)	30(b)	43	43	30(b)	30(b)	30(b)	30(b)	30(b)	30(b)	
January ..	29.865	6.8	19.7	23/47	58	SE	NE	6.74	9.8	5.7	3.5
February ..	29.912	7.0	23.2	21/54	67	SE	NE	5.49	6.5	5.6	2.4
March ..	29.975	6.5	20.3	1/29	65	S	E	5.05	5.9	5.1	5.4
April ..	30.035	5.9	16.7	3/25	57	S	E	4.05	5.0	4.3	7.8
May ..	30.083	5.8	17.9	17/26	49	SW	SE	3.09	4.1	4.3	8.3
June ..	30.091	5.7	19.0	14/28	58	SW	W & SW	2.45	2.9	4.4	9.2
July ..	30.090	5.6	22.0	13/54	52	SW	W & SW	2.69	2.8	3.8	12.4
August ..	30.105	5.8	14.8	4/35	56	SW	NE	3.51	3.8	3.1	13.1
September ..	30.067	5.9	16.1	1/48	57	SW	NE	4.51	5.8	3.3	13.0
October ..	30.019	6.3	15.7	1/41	62	S	NE	5.81	7.1	4.2	8.5
November ..	29.958	6.7	15.5	10/28	62	SE & N	NE	6.32	9.5	4.9	5.9
December ..	29.890	7.0	19.5	15/26	79	SE	NE	7.02	10.6	5.3	3.8
Year { Totals ..	—	—	—	—	—	—	—	56.73	73.8	—	93.3
Year { Averages ..	30.007	6.3	—	—	—	SW	NE	—	—	—	—
Year { Extremes ..	—	—	23.2	21/2/54	79	—	—	—	—	—	—

(a) Scale 0-10. (b) Standard 30 years' normal (1911-1940).

Temperature and Sunshine.

Month.	Mean Temperature (°Fahr.).			Extreme Shade Temperature (°Fahr.).		Extreme Range.	Extreme Temperature (°Fahr.).		Mean Daily Hours of Sunshine.				
	Mean Max.	Mean Min.	Mean	Highest.	Lowest.		Highest in Sun.	Lowest on Grass.					
										71		71	
No. of years over which observation extends.	30(a)	30(a)	30(a)	71	71	71	50(b)	71	30(a)				
January ..	85.5	69.1	77.3	109.8	26/40	58.8	4/93	51.0	169.0	2/37	49.9	4/93	7.6
February ..	84.6	68.7	76.6	105.7	21/25	58.5	23/31	47.2	165.2	6/10	49.1	22/31	7.4
March ..	82.3	66.2	74.3	99.4	5/19	52.4	29/13	47.0	162.5	6/39	45.4	29/13	7.0
April ..	79.1	61.5	70.3	95.2	(c)	44.4	25/25	50.8	153.8	11/16	36.7	24/25	7.1
May ..	73.7	55.6	64.7	90.3	21/23	40.6	30/51	49.7	147.0	1/10	29.8	8/97	6.6
June ..	69.4	51.5	60.5	88.9	19/18	36.3	29/08	52.6	136.0	3/18	25.4	23/88	6.3
July ..	68.6	49.4	59.0	84.3	23/46	36.1	(d)	48.2	146.1	20/15	23.9	11/90	6.8
August ..	71.1	50.0	60.6	91.0	14/46	37.4	6/87	53.6	141.9	20/17	27.1	9/99	7.9
September ..	75.5	54.8	65.1	100.9	22/43	40.7	1/96	60.2	155.5	26/03	30.4	1/89	8.2
October ..	79.2	60.3	69.8	101.4	18/93	43.3	3/99	58.1	157.4	31/18	34.9	8/89	8.4
November ..	82.3	64.6	73.4	106.1	18/13	48.5	2/05	57.6	162.3	7/89	38.8	1/05	8.2
December ..	84.5	67.5	76.0	105.9	26/93	56.4	13/12	49.5	165.9	28/42	49.1	3/94	8.2
Year { Averages ..	78.0	59.9	69.0	—	—	—	—	—	—	—	—	—	7.5
Year { Extremes ..	—	—	—	109.8	26/1/40	36.1	(d)	73.7	169.0	2/11/37	23.9	11/7/90	—

(a) Standard 30 years' normal (1911-1940). (b) From 1887 to March, 1947, excluding 1927 to 1936.
(c) 9/1896 and 5/1903. (d) 12/7/1894 and 2/7/1896.

Humidity, Rainfall and Fog.

Month.	Vapour Pressure (inches)	Rel. Hum. (%) at 9 a.m.			Rainfall (inches).					Fog. Mean No. of Days of Fog.				
		Mean 9 a.m.	Mean.	Highest Mean.	Lowest Mean.	Mean Monthly.	Mean No. of Days of Rain.	Greatest Monthly.	Least Monthly.		Greatest in One Day.			
												106		106(b)
No. of years over which observation extends.	30(a)	30(a)	69	69	30(a)	30(a)	106	106(b)	106	30(a)				
January ..	0.636	66	79	53	5.72	12	27.72	1895	0.32	1919	18.31	21/87	0.6	
February ..	0.644	69	82	55	5.47	12	40.39	1893	0.58	1849	10.61	6/31	0.9	
March ..	0.606	72	85	56	4.97	14	34.04	1870	Nil	1849	11.18	14/08	1.6	
April ..	0.512	71	80	56	3.68	11	15.28	1867	0.04	1944	5.46	5/33	4.0	
May ..	0.420	71	85	59	2.35	9	13.85	1867	Nil	1846	5.62	9/79	5.4	
June ..	0.357	73	84	54	2.75	8	14.03	1873	Nil	1847	6.41	15/48	4.5	
July ..	0.331	71	88	53	1.88	8	8.60	1950	Nil	1841	3.54	(c)	4.9	
August ..	0.338	67	80	53	1.07	7	14.67	1879	Nil	(d)	4.89	12/87	5.9	
September ..	0.396	62	76	47	1.69	7	5.43	1886	0.10	1907	2.46	2/94	2.8	
October ..	0.459	59	72	48	2.27	8	11.41	1949	0.03	1948	5.34	25/49	4.0	
November ..	0.533	61	73	45	4.00	10	12.40	1917	Nil	1842	4.46	16/86	0.7	
December ..	0.589	62	70	51	4.24	11	17.36	1942	0.35	1865	6.60	28/71	0.4	
Year { Totals ..	—	—	—	—	40.09	117	—	—	—	—	—	—	—	33.3
Year { Averages ..	0.485	67	—	—	—	—	—	—	—	—	—	—	—	—
Year { Extremes ..	—	—	88	45	—	—	40.39	2/1893	Nil	(e)	18.31	21/11/87	—	

(a) Standard 30 years' normal (1911-1940). (b) Records incomplete for various years between 1846 and 1859. (c) 15/1876 and 16/1889. (d) 1862, 1869, 1880. (e) Various months in various years.

CLIMATOLOGICAL DATA: SYDNEY, NEW SOUTH WALES.
(LAT. 33° 52' S., LONG. 151° 12' E. HEIGHT ABOVE M.S.L. 138 FT.)
Barometer, Wind, Evaporation, Lightning, Clouds and Clear Days.

Month.	Bar. corrected to 12° F. Mm. Sea Level and Standard Gravity from 9 a.m. and 3 p.m. readings.	Wind. (Height of Anemometer 58 feet.)						Mean Amount of Evaporation (inches).	No. of Days of Lightning.	Mean Amount of Clouds, 9 a.m. to 3 p.m., 9 p.m. (g)	No. of Clear Days.
		Average Miles per Hour.	Highest Mean Speed in One Day (miles per hour).	Highest Gust Speed (miles per hour).	Prevailing Direction.		26(c)				
					9 a.m.	3 p.m.					
No. of years of observations.	30(b)	26(c)	44(d)	38(e)	26(c)	26(c)	26(c)	30(f)	30(b)	30(b)	
January ..	29.875	8.9	24.9	2/22	74	S	ENE	5.71	4.8	5.7	4.8
February ..	29.942	8.1	23.5	19/57	63	NE	ENE	4.68	3.3	5.5	4.4
March ..	30.009	7.5	20.7	10/44	58	W	ENE	4.05	2.8	5.3	3.8
April ..	30.063	7.0	23.4	19/27	72	W	NE	2.91	2.4	5.0	3.4
May ..	30.098	6.8	19.6	2/26	63	W	S	2.17	1.6	4.9	7.0
June ..	30.078	7.1	24.5	17/14	73	W	W	1.61	1.5	4.8	8.3
July ..	30.070	7.2	26.6	6/31	68	W	W	1.69	1.1	4.9	10.1
August ..	30.060	7.4	24.6	9/51	68	W	NE	2.30	2.1	3.9	11.1
September ..	30.018	8.0	22.3	19/17	70	W	ENE	4.17	3.9	5.0	10.4
October ..	29.976	8.2	23.3	2/57	95	W	ENE	4.97	4.5	5.5	5.7
November ..	29.935	8.5	22.6	14/30	71	W & E	ENE	4.17	5.4	5.8	5.7
December ..	29.881	8.9	24.9	10/20	75	S	ENE	5.64	5.4	5.8	4.8
Year { Totals ..	30.000	7.8	—	—	—	W	NE	42.90	36.4	—	87.8
{ Averages ..	—	—	—	—	—	—	—	—	—	5.0	—
{ Extremes ..	—	—	26.6	6/7/31	95	—	—	—	—	—	—

(a) Scale 0-10. (b) Standard 30 years' normal (1911-1940). (c) 1915-1940. (d) 1914-1953. (e) 1917-1954. (f) 1921-1950.

Temperature and Sunshine.

Month.	Mean Temperature (°Fahr.).			Extreme Shade Temperature (°Fahr.).		Extreme Range.	Extreme Temperature (°Fahr.).		Mean Daily Hours of Sunshine.				
	Mean Max.	Mean Min.	Mean	Highest.	Lowest.		Highest in Sun.	Lowest on Grass.					
										30(a)	30(a)	30(a)	99
No. of years over which observation extends.	30(a)	30(a)	30(a)	99	99	99	84	99	30(b)				
January ..	78.6	65.1	71.8	113.6	14/39	51.1	18/49	62.5	164.3	26/15	43.7	6/25	7.5
February ..	78.7	65.5	72.1	107.8	8/26	49.3	28/63	58.5	168.3	14/39	42.8	22/33	7.0
March ..	76.6	62.9	69.8	102.6	3/69	48.8	14/86	53.8	158.3	10/26	39.9	17/13	6.4
April ..	72.0	57.7	64.9	91.4	1/36	44.6	27/64	46.8	144.1	10/77	33.3	24/09	6.1
May ..	67.0	52.4	59.7	86.0	1/19	40.2	22/59	45.8	129.7	1/96	29.3	25/17	5.7
June ..	62.8	48.1	55.5	80.4	11/31	35.7	22/32	44.7	125.5	2/23	28.0	22/32	5.3
July ..	61.8	46.4	54.1	78.3	22/26	35.9	12/90	42.4	124.7	19/77	24.0	4/93	6.1
August ..	64.3	47.6	56.0	82.8	12/46	36.8	3/72	46.0	149.0	30/78	26.1	4/09	7.0
September ..	68.3	51.4	59.9	92.3	27/19	40.8	2/45	51.5	142.2	12/78	30.1	17/05	7.3
October ..	71.7	55.9	63.8	99.4	4/42	42.2	6/27	57.2	152.2	20/33	32.7	9/05	7.5
November ..	74.5	59.8	67.1	104.5	6/46	45.8	1/05	61.3	158.5	28/99	36.0	6/06	7.5
December ..	76.9	63.2	70.1	108.0	20/57	48.4	3/24	59.6	164.5	27/89	41.4	3/24	7.5
Year { Averages ..	71.1	56.3	63.7	—	—	—	—	—	—	—	—	—	6.8
{ Extremes ..	—	—	—	113.6	14/11/39	35.7	22/6/32	77.9	168.3	14/2/39	24.0	4/7/93	—

(a) Standard 30 years' normal (1911-1940). (b) 1921-1950 (different exposure prior to 1921).

Humidity, Rainfall and Fog.

Month.	Vapour Pressure (inches).	Rel. Hum. (%) at 9 a.m.				Rainfall (inches).				Fog. Mean No. of Days of Fog.			
		Mean 9 a.m.	Mean.	Highest Mean.	Lowest Mean.	Mean Monthly.	Mean No. of Days of Rain.	Greatest Monthly.	Least Monthly.		Greatest in One Day.		
												30(a)	30(a)
No. of years over which observation extends.	30(a)	30(a)	80	80	30(a)	30(a)	99	99	99	30(b)			
January ..	0.537	65	78	58	3.86	13	15.26	1911	0.25	1932	7.08	13/11	0.4
February ..	0.560	68	81	60	3.15	12	22.22	1956	0.12	1939	8.90	25/73	0.8
March ..	0.527	71	85	62	4.44	13	20.52	1942	0.42	1876	11.05	28/42	1.8
April ..	0.441	73	87	63	5.65	14	24.49	1861	0.06	1868	7.52	29/60	2.8
May ..	0.362	75	90	63	4.98	12	23.03	1919	0.14	1957	8.36	28/89	2.7
June ..	0.303	76	89	63	3.68	11	25.30	1950	0.19	1904	5.17	16/84	3.3
July ..	0.282	74	88	63	4.89	12	13.23	1950	0.10	1946	7.80	7/31	2.9
August ..	0.288	68	84	54	2.41	10	14.89	1899	0.04	1885	5.33	2/60	2.3
September ..	0.325	62	79	49	2.77	11	14.05	1879	0.08	1882	5.69	10/79	1.0
October ..	0.378	60	77	46	2.80	11	11.13	1916	0.21	1867	6.37	13/02	0.6
November ..	0.433	60	79	42	2.54	11	9.88	1865	0.07	1915	4.23	19/00	0.6
December ..	0.501	63	77	51	3.63	13	15.82	1920	0.23	1913	4.75	13/10	0.4
Year { Totals ..	—	—	—	—	44.80	143	—	—	—	—	—	—	20.6
{ Averages ..	0.393	68	—	—	—	—	—	—	—	—	—	—	—
{ Extremes ..	—	—	90	42	—	—	25.30	6/1950	0.04	8/1885	11.05	28/3/42	—

(a) Standard 30 years' normal (1911-1940). (b) 1921-1950.

CLIMATOLOGICAL DATA; CANBERRA, AUSTRALIAN CAPITAL TERRITORY.

(LAT. 35° 18' S., LONG. 149° 6' E. HEIGHT ABOVE M.S.L., 1,906 FT.)

Barometer, Wind, Evaporation, Lightning, Clouds and Clear Days.

Month.	Bar. corrected to 32° F. Min. Sea level and Standard Gravity from 9 a.m. and 3 p.m. readings.	Wind. (Height of Anemometer 20 feet.)					Mean Amount of Evaporation (inches).	No. of Days of Lightning.	Mean Amount of Clouds 9 a.m. and 3 p.m. (e)	No. of Clear Days.	
		Average Miles per Hour.	Highest Mean Speed in One Day (miles per hour).	Highest Gust Speed (miles per hour).	Prevailing Direction.						
					9 a.m.	3 p.m.					
No. of years of observations.	26	27	28	(b)	27	27	28	20	26	27	
January ..	29.856	4.7	14.9	23/33	—	NW	NW	8.31	1.5	4.9	7.3
February ..	29.900	4.2	15.3	24/33	—	E	NW	6.42	2.3	5.1	6.3
March ..	30.009	3.7	18.2	28/42	—	—	NW	5.20	0.2	5.1	6.9
April ..	30.059	3.6	18.6	8/45	—	NW	NW	3.28	0.3	5.4	4.7
May ..	30.126	3.0	12.6	3/30	—	NW	NW	1.95	0.2	5.6	5.8
June ..	30.120	3.6	16.1	7/31	—	NW	NW	1.29	0.1	6.0	5.3
July ..	30.133	3.4	23.4	7/31	—	NW	NW	1.27	0.0	5.7	5.6
August ..	30.065	4.1	15.7	25/36	—	NW	NW	1.81	0.1	5.4	5.7
September ..	30.057	4.2	17.4	28/34	—	NW	NW	2.87	0.4	5.1	6.1
October ..	29.954	4.3	14.7	12/57	—	NW	NW	4.43	1.0	5.4	5.2
November ..	29.885	4.7	17.2	28/42	—	NW	NW	5.87	1.1	5.5	4.5
December ..	29.834	4.7	16.1	11/38	—	NW	NW	7.64	0.7	5.0	6.3
Year { Totals ..	30.000	4.0	—	—	—	NW	NW	50.34	7.9	—	68.9
{ Averages ..	—	—	—	23.4 7/7/31	—	—	—	—	—	—	—
{ Extremes ..	—	—	—	—	—	—	—	—	—	—	—

(a) Scale 0-10.

(b) No record.

Temperature and Sunshine.

Month.	Mean Temperature (°Fahr.).			Extreme Shade Temperature (°Fahr.).		Extreme Range.	Extreme Temperature (°Fahr.).		Mean Daily Hours of Sunshine.				
	Mean.	Min.	Mean.	Highest.	Lowest.		Highest in Sun.	Lowest on Grass.					
										30	30	(a)	30
No. of years over which observation extends.	29	29	29	30	30	30	—	—	27				
January ..	82.4	56.0	69.2	107.4	11/39	38.0	1/56	69.4	—	30.1	10/50	8.4	
February ..	80.7	56.1	68.4	99.8	13/33	35.0	(b)	64.8	—	26.5	23/43	7.3	
March ..	76.2	52.7	64.4	99.1	6/38	34.8	31/49	64.3	—	26.4	26/35	7.2	
April ..	66.7	45.5	56.1	89.7	6/38	29.0	29/34	60.7	—	19.0	18/44	6.7	
May ..	59.3	39.1	49.2	72.6	1/36	22.5	(c)	50.1	—	15.6	(d)	5.2	
June ..	52.6	35.7	44.1	64.9	1/54	18.1	20/35	43.9	—	8.9	25/44	4.2	
July ..	51.8	33.8	42.8	63.5	16/34	20.0	(e)	43.5	—	10.8	9/37	4.8	
August ..	55.1	35.4	45.3	71.0	24/54	21.0	3/29	49.5	—	10.1	6/44	5.8	
September ..	61.4	38.9	50.1	81.5	16/34	25.2	6/46	56.3	—	13.0	6/45	7.2	
October ..	67.0	44.2	55.6	90.0	13/46	29.0	24/28	61.0	—	18.2	2/45	7.8	
November ..	72.9	48.7	60.8	101.4	19/44	32.2	11/36	69.2	—	22.9	6/56	8.2	
December ..	79.5	53.3	66.4	103.5	27/38	36.0	24/28	67.5	—	29.1	21/56	8.5	
Year { Averages ..	67.1	44.9	56.0	—	—	—	—	—	—	—	—	6.8	
{ Extremes ..	—	—	—	107.4	11/11/39	18.1	20/6/35	89.3	—	—	8.9	25/6/44	—

(a) No record. 9/37 and 27/43.

(b) 22/31 and 23/31.

(c) 9/29 and 25/57.

(d) 13/37 and 15/46.

(e) 19/29.

Humidity, Rainfall and Fog.

Month.	Vapour Pressure (inches).	Rel. Hum. (%) at 9 a.m.			Rainfall (inches).				Fog. Mean No. of Days of Fog.				
		Mean.	Highest.	Lowest.	Mean Monthly.	Mean No. of Days of Rain.	Greatest Monthly.	Least Monthly.		Greatest in One Day.			
											28	28	28
No. of years over which observation extends.	26	28	28	28	29	29	30	30	30	25			
January ..	0.370	53	69	39	2.13	7	6.69	1941	0.02	1932	2.47	19/50	0.1
February ..	0.388	59	71	40	2.17	7	6.03	1948	0.01	1933	3.24	17/28	0.2
March ..	0.378	66	82	48	2.48	7	12.69	1950	0.01	1940	2.53	20/52	1.0
April ..	0.315	71	81	54	2.17	8	5.19	1952	0.07	1942	2.52	9/45	1.4
May ..	0.254	79	89	67	2.06	8	6.13	1948	0.06	1935	3.88	3/48	4.8
June ..	0.212	81	90	72	1.92	9	6.09	1931	0.18	1944	2.32	25/56	5.8
July ..	0.196	81	91	73	1.61	10	4.09	1933	0.27	1940	2.02	13/33	5.9
August ..	0.213	75	88	60	1.98	11	4.71	1939	0.36	(a)	2.07	12/29	2.4
September ..	0.239	66	78	51	1.62	9	3.03	1937	0.13	1946	1.75	3/47	1.4
October ..	0.273	60	72	46	2.77	11	6.59	1934	0.34	1940	2.51	25/34	0.4
November ..	0.301	55	67	38	2.11	8	4.45	1950	0.28	1936	2.45	9/50	0.1
December ..	0.338	51	70	37	1.86	8	8.80	1947	0.16	1938	2.29	28/29	0.0
Year { Totals ..	—	—	—	—	24.88	103	—	—	—	—	—	—	22.9
{ Averages ..	0.286	66	—	—	—	—	—	—	—	—	—	—	—
{ Extremes ..	—	—	91	37	—	—	12.69	3/50	0.01	2/33,3/40	3.88	3/5/48	—

(a) 1944 and 1949.

CLIMATOLOGICAL DATA; MELBOURNE, VICTORIA.
(LAT. 37° 49' S., LONG. 144° 58' E. HEIGHT ABOVE M.S.L. 114 FT.)
Barometer, Wind, Evaporation, Lightning, Clouds and Clear Days.

Month.	Bar. corrected to 32° F. Ven. Sea Level and Standard Gravity from 9 a.m. and 3 p.m. readings.	Wind. (Height of Anemometer 93 feet.)					Mean Amount of Evaporation (inches).	No. of Days of Lightning.	Mean Amount of Clouds, 9 a.m., 3 p.m., 9 p.m.(g)	No. of Clear Days.	
		Average Miles per Hour.	Highest Mean Speed in One Day (miles per hour).	Highest Gust Speed (miles per hour).	Prevailing Direction.						
					9 a.m.	3 p.m.					
No. of years of observations.	30(b)	15(c)	45	48	30(b)	30(b)	30(b)	30(b)	30(b)		
January	29.897	8.8	21.1	27/41	66	S & SW	S	6.55	1.8	4.9	6.8
February	29.950	8.4	19.0	13/47	74	N & S	S	5.10	2.3	4.8	6.4
March	30.025	7.8	17.2	19/50	66	N	S	4.26	1.8	5.3	5.5
April	30.092	7.1	19.9	16/43	67	N	S	2.53	1.2	5.9	4.6
May	30.113	7.4	21.8	1/57	72	N	S	1.57	0.5	6.1	3.4
June	30.097	7.2	22.8	16/47	62	N	S	1.18	0.4	6.5	2.7
July	30.079	8.7	20.9	9/44	68	N	S	1.16	0.3	6.3	2.9
August	30.048	8.2	21.3	20/42	65	N	S	1.54	0.9	6.0	3.1
September	30.001	8.5	20.9	15/57	69	N & W	N & S	2.41	1.3	5.9	3.5
October	29.968	8.4	18.6	12/52	69	N	S	3.54	1.8	6.1	3.8
November	29.951	8.6	19.4	4/50	71	S & SW	S	4.62	2.3	6.0	3.6
December	29.896	8.7	21.0	11/52	61	S & SW	S	5.85	1.9	5.6	4.5
Year { Totals	—	—	—	—	—	—	—	40.31	16.5	—	50.6
Year { Averages	30.010	8.1	—	—	—	N	S	—	—	—	—
Year { Extremes	—	—	22.8	16/6/47	74	—	—	—	—	—	—

(a) Scale 0-10. (b) Standard 30 years' normal (1911-1940). (c) Early records not comparable.

Temperature and Sunshine.

Month.	Mean Temperature (°Fahr.).			Extreme Shade Temperature (°Fahr.).		Extreme Range.	Extreme Temperature (°Fahr.).		Mean Daily Hours of Sunshine.				
	Mean Max.	Mean Min.	Mean	Highest.	Lowest.		Highest in Sun.	Lowest on Grass.					
										No. of years over which observation extends.			
No. of years over which observation extends.	30(a)	30(a)	30(a)	102	102	102	86(b)	98	35(c)				
January	77.7	56.9	67.3	114.1	13/39	42.0	28/85	72.1	178.5	14/62	30.2	28/85	7.8
February	78.6	58.0	68.3	109.5	7/01	40.2	24/24	69.3	167.5	15/70	30.9	6/91	7.4
March	74.9	55.2	65.1	107.0	11/40	37.1	17/84	69.9	164.5	1/68	28.9	(d)	6.5
April	67.9	50.8	59.3	94.8	5/38	34.8	24/88	60.0	152.0	8/61	25.0	23/97	5.0
May	62.0	46.9	54.5	83.7	7/05	29.9	29/116	53.8	142.6	2/59	21.1	26/16	4.1
June	56.8	43.8	50.3	72.3	2/57	28.0	11/66	44.3	129.0	11/61	19.9	30/29	3.4
July	56.2	42.6	49.4	69.3	22/26	27.0	21/69	42.3	125.8	27/80	20.5	12/03	3.7
August	58.7	43.7	51.2	77.0	20/85	28.3	11/63	48.7	137.4	29/69	21.3	14/02	4.6
September	63.3	46.0	54.7	88.6	28/28	31.0	3/40	57.6	142.1	20/67	22.8	8/18	5.5
October	67.9	48.7	58.3	98.4	24/14	32.1	3/71	66.3	154.3	28/68	24.8	22/18	5.8
November	71.3	51.8	61.5	105.7	27/94	36.5	2/96	69.2	159.6	29/65	24.6	2/96	6.2
December	75.4	55.3	65.3	110.7	15/76	40.0	4/70	70.7	170.3	20/69	33.2	1/04	7.0
Year { Averages	67.6	50.0	58.8	—	—	—	—	—	—	—	—	—	—
Year { Extremes	—	—	—	114.1	13/11/39	27.0	21/7/69	87.1	178.5	14/11/62	19.9	30/6/29	—

(a) Standard 30 years' normal (1911-1940). (b) Records discontinued, 1946. (c) 1916-1950. (d) 17/1884 and 20/1897.

Humidity, Rainfall and Fog.

Month.	Vapour Pressure (inches)	Rel. Hum. (%) at 9 a.m.			Rainfall (inches).					Fog. Mean No. of Days of Fog.			
		Mean 9 a.m.	Highest Mean.	Lowest Mean.	Mean Monthly.	Mean No. of Days of Rain.	Greatest Monthly.	Least Monthly.	Greatest in One Day.				
											No. of years over which observation extends.		
No. of years over which observation extends.	30(a)	30(a)	48	48	30(a)	30(a)	102	102	102	30(a)			
January	0.382	58	68	50	1.88	9	6.66	1941	0.01	1932	2.97	9/97	0.1
February	0.417	62	77	48	2.00	8	7.72	1939	0.03	1870	3.44	26/46	0.3
March	0.385	64	79	50	2.22	9	7.50	1911	0.14	1934	3.55	5/19	1.1
April	0.351	72	82	66	2.30	13	6.71	1901	Nil	1923	2.28	22/01	2.3
May	0.311	79	88	70	1.94	14	5.60	1942	0.14	1934	1.85	7/91	6.8
June	0.276	83	92	75	2.06	16	4.51	1859	0.73	1877	1.74	21/04	6.5
July	0.264	82	86	75	1.93	17	7.02	1891	0.57	1902	2.71	12/91	6.5
August	0.271	76	82	65	2.02	17	4.35	1939	0.48	1903	1.94	26/24	3.7
September	0.288	68	76	60	2.20	15	7.93	1916	0.52	1907	2.62	12/80	1.3
October	0.307	62	67	52	2.63	14	7.61	1869	0.29	1914	3.00	17/69	0.3
November	0.336	60	69	52	2.33	13	8.11	1954	0.25	1895	2.86	21/54	0.3
December	0.373	59	69	48	2.38	11	7.18	1863	0.11	1904	3.92	4/54	0.2
Year { Totals	—	—	—	—	25.89	156	—	—	—	—	—	—	—
Year { Averages	0.323	69	—	—	—	—	—	—	—	—	—	—	—
Year { Extremes	—	—	92	48	—	—	8.11	11/1954	Nil	4/1923	3.92	4/12/54	—

(a) Standard 30 years' normal (1911-1940).

CLIMATOLOGICAL DATA; HOBART, TASMANIA.
(LAT. 42° 53' S., LONG. 147° 20' E. HEIGHT ABOVE M.S.L. 177 FT.)
Barometer, Wind, Evaporation, Lightning, Clouds and Clear Days.

Month.	Bar. corrected to 32° F. Mm. Sea Level and Standard Gravity from 9 a.m. and 3 p.m. readings.	Wind.					Mean Amount of Evaporation (inches).	No. of Days of Lightning.	Mean Amount of Clouds, 9 a.m., 3 p.m., 9 p.m.(g)	No. of Clear Days.	
		(Height of Anemometer 40 feet.)									
		Average Miles per Hour.	Highest Mean Speed in One Day (miles per hour).	Highest Gust Speed (miles per hour).	Prevailing Direction.						
No. of years of observations.	30(b)	30(b)	65	65	9 a.m.	3 p.m.	30(b)	30(b)	30(b)		
January ..	29.819	8.0	20.8	30/16	76	NNW	SSE	4.84	0.9	6.4	1.9
February ..	29.913	7.2	25.2	4/27	65	NNW	SSE	3.71	1.0	6.2	2.3
March ..	29.961	6.8	21.4	13/38	75	NW	SSE	3.10	1.2	6.1	2.4
April ..	29.997	6.7	24.1	9/52	74	NW	W	1.98	0.7	6.5	1.7
May ..	30.009	6.3	20.2	20/36	79	NNW	NW	1.37	0.4	6.1	2.4
June ..	29.986	6.2	23.7	27/20	71	NW	NW	0.91	0.4	6.2	2.4
July ..	29.958	6.5	22.9	22/53	78	NNW	NNW	0.94	0.3	6.1	2.0
August ..	29.906	6.8	25.5	19/26	87	NNW	NW	1.28	0.4	6.1	2.1
September ..	29.860	7.9	21.5	26/15	84	NNW	NW	1.97	0.7	6.3	1.5
October ..	29.833	8.2	19.2	8/12	74	NNW	SW	3.05	0.6	6.6	1.0
November ..	29.831	7.9	21.2	18/15	73	NNW	S	3.77	0.7	6.4	1.3
December ..	29.816	7.6	23.4	1/34	70	NNW	SSE	4.37	0.5	6.8	1.1
Year { Totals ..	29.907	7.2	—	—	—	NNW	W	31.29	7.8	—	22.1
Year { Averages ..	—	—	—	—	—	—	—	—	—	—	—
Year { Extremes ..	—	—	25.5	19/8/26	87	—	—	—	—	—	—

(a) Scale 0-10. (b) Standard 30 years' normal (1911-1940).

Temperature and Sunshine.

Month.	Mean Temperature (°Fahr.).			Extreme Shade Temperature (°Fahr.).		Extreme Range.	Extreme Temperature (°Fahr.).		Mean Daily Hours of Sunshine.				
	Mean Max.	Mean Min.	Mean	Highest.	Lowest.		Highest in Sun.	Lowest on Grass.					
										30(a)	30(a)	30(a)	88(b)
No. of years over which observation extends.	30(a)	30(a)	30(a)	88(b)	88(b)	88(b)	57(c)	70(b)	30				
January ..	69.8	52.4	61.0	105.0	1/00	40.1 (d)	64.9	160.0 (e)	30.6	19/97	7.7		
February ..	70.6	53.7	62.2	104.4	12/99	39.0	20/87	65.4	165.0	24/98	28.3	—/87	7.1
March ..	67.5	51.3	59.4	99.1	13/40	35.2	31/26	63.9	150.9	26/44	27.5	30/02	6.4
April ..	62.2	48.0	55.1	87.1	1/41	33.3	24/88	53.8	142.0	18/93	25.0	—/86	5.0
May ..	57.8	44.6	51.2	77.8	5/21	29.2	20/02	48.6	128.0 (f)	20.0	19/02	4.4	
June ..	52.8	41.2	47.0	69.2	1/07	29.2	28/44	40.0	122.0	12/94	21.0	6/87	4.0
July ..	52.7	40.6	46.6	66.1	14/34	27.7	11/95	38.4	121.0	12/93	18.7	16/86	4.4
August ..	55.4	41.7	48.7	71.6	28/14	28.9	9/51	42.7	129.0	—/87	20.1	7/09	5.1
September ..	59.0	43.7	51.4	81.7	23/26	31.0	16/97	50.7	138.0	23/93	18.3	16/26	5.9
October ..	62.5	46.1	54.3	92.0	24/14	32.0	12/89	60.0	156.0	9/93	23.8	(g)	6.1
November ..	65.0	48.2	56.6	98.3	26/37	35.0	16/41	63.3	154.0	19/92	26.0	1/08	7.2
December ..	67.9	51.3	59.6	105.2	30/97	38.0	3/06	67.2	161.5	10/39	27.2	—/86	7.3
Year { Averages ..	61.9	46.9	54.4	—	—	—	—	—	—	—	—	—	5.9
Year { Extremes ..	—	—	—	105.2	30/12/97	27.7	11/7/95	77.5	165.0	24/2/98	18.3	16/9/26	—

(a) Standard 30 years' normal (1911-1940). (b) Records 1855-1882 not comparable. (c) Period 1934-1938 not comparable; records discontinued, 1946. (d) 9/37 and 11/37. (e) 5/86 and 13/05. (f) —/89 and —/93. (g) 1/86 and —/99.

Humidity, Rainfall and Fog.

Month.	Vapour Pressure (inches)	Rel. Hum. (%) at 9 a.m.			Rainfall (inches).					Fog. Mean No. of Days of Fog.			
		Mean.	Highest Mean.	Lowest Mean.	Mean Monthly.	Mean No. of Days of Rain.	Greatest Monthly.	Least Monthly.	Greatest in One Day.				
											30(a)	30(a)	75(b)
No. of years over which observation extends.	30(a)	54	69	69	30(a)	30(a)	75(b)	75(b)	75(b)	30(c)			
January ..	0.309	59	72	46	1.82	13	5.91	1893	0.17	1915	2.96	30/16	0.0
February ..	0.342	63	77	48	1.68	10	5.01	1956	0.11	1914	2.20	1/54	0.0
March ..	0.323	67	77	52	2.13	13	10.05	1946	0.29	1943	3.47	17/46	0.3
April ..	0.290	72	84	58	2.31	14	8.50	1935	0.07	1904	5.02	20/09	0.2
May ..	0.263	78	89	65	1.71	14	6.37	1905	0.14	1913	1.75	2/93	0.9
June ..	0.233	80	91	65	2.25	16	8.15	1889	0.28	1886	5.80	7/54	0.8
July ..	0.227	80	94	72	2.14	17	6.02	1922	0.17	1950	2.51	18/22	1.0
August ..	0.232	76	92	60	1.82	18	6.32	1946	0.30	1892	2.28	14/90	0.4
September ..	0.240	67	85	58	1.90	17	7.93	1957	0.38	1951	6.15	15/57	0.1
October ..	0.258	63	73	51	2.52	18	7.60	1947	0.39	1914	2.58	4/06	0.0
November ..	0.274	60	72	50	2.23	16	7.39	1885	0.33	1921	3.70	30/85	0.1
December ..	0.306	58	67	45	2.52	14	7.72	1916	0.17	1931	3.33	5/41	0.0
Year { Totals ..	—	—	—	—	25.03	180	—	—	—	—	—	—	3.8
Year { Averages ..	0.271	69	—	—	—	—	—	—	—	—	—	—	—
Year { Extremes ..	—	—	94	45	—	—	10.05	3/1946	0.07	4/1904	5.80	7/6/54	—

(a) Standard 30 years' normal (1911-1940). (b) Records prior to 1883 not comparable. (c) 1922-1951.